

Evaluation of Different Techniques for Breaking Seed Dormancy of *Heliotropium europaeum* L. (Boraginaceae)

Ali Asghar Aliloo* and Shabnam Darabinejad

Department of Agronomy and Plant Breeding, Faculty of Agriculture, University of Maragheh, Maragheh, P.O. Box 55181-83111, IRAN

Received: 16.07.2013; Accepted: 22.07.2013; Published Online: 29.08.2013

ABSTRACT

Prediction of germination and emergence of weeds plays a key role in weed management strategies. For this reason, the seed dormancy status of *Heliotropium europaeum* (Boraginaceae) was investigated in North West of Iran. The dormancy breaking treatments were after-ripening, gibberellic acid, hot water, KNO₃, cold stratification, and washing. Seeds were strongly dormant at maturity and maintained high levels of dormancy for long periods of time. However, after 12 months storage at dry conditions, germination percentage increased to 37%. Washing and hot water treatments did not alter seed dormancy behavior, while the influence of gibberellic acid, KNO₃ and cold stratification was significant. The highest value for germination percentage obtained by cold stratification increased the value up to 90%. Results of seedling growth showed that compared to other treatments, the seeds treated by KNO₃ produced longer shoots and vigorous seedlings. The high vigor index was also recorded for cold stratification. It is concluded that the seeds of *Heliotrope* had a high primary dormancy and among treatments, the cold stratification was a more effective method in breaking seed dormancy .

Key Words: Dormancy, heliotrope, germination, seedling growth

INTRODUCTION

Heliotropium europaeum L., belonging to the Boraginaceae family, is a summer annual weed, which successfully grows on road margins, bare and cultivated fields in North West of Iran (Azerbaijan). Seeds germinate in the warm, moist conditions after late spring or early summer. In rotation crop systems, the fallow year provides a condition in which the population of this plant increases at a high rate because *Heliotrope* has a massive seeding potential and the seeds are viable for many years in the soil seed bank. The plant contains toxins (Pyrrolizidine alkaloids) that exist in its all parts (Saedi and Semnani 2009). It can be used for medicinal purposes to treat warts, stimulate bile secretion, regulate menstruation, lower fever, and soothe insect bites, gout, and inflammation in joints.

During seed development, the embryo gradually enters a quiescent phase due to desiccation. After dispersal from the mother plant, seeds may germinate under suitable conditions, which are specific for each species. However, in many circumstances the germination of a viable seed will not take place even though all the necessary environmental conditions for embryo growth are satisfied. This phenomenon is called seed dormancy. Seed dormancy brings some advantages to the plant. For instance, it provides enough time for seed dispersal over greater distances. It also maximizes seedling survival by the timing of germination under unfavorable conditions. Several external factors release the seed from dormancy, and dormant seeds typically respond to more than one condition. A common method used to release dormancy is after-ripening which is a period of dry storage at room temperature of freshly harvested, mature seeds (Finch-Savage and Leubner-Metzger 2006). Another useful method to overcome dormancy in hard seeds is the hot water. This treatment cracks the macrosclerid layer or affects the strophilar plug (Francisco *et al.* 2001). This method is also effectively used to break dormancy in rice seeds (Tung *et al.* 2011) and Orka (Mohammadi *et al.* 2012). A stimulatory effect on germination of *Commelina benghalensis* L. seeds has been also reported by hot water at 70°C (Kim *et al.* 1990). Soaking of *Nicandra physalodes* L. Seeds in preheated water at 50°C for 10 minutes and incubated in the light at 25°C did not affect the germination, but the same treatment followed by incubation in the dark instead of light significantly reduced the germination (Watanabe *et al.* 2002). In some species, the accumulation of germination

* Corresponding author: aliloo@maragheh.ac.ir

inhibitors causes suppression on germination. If they are water soluble, repeated washing with water or soaking in tap water can remove chemical inhibitors and release their dormancy (Bewley and Black 1994). Abscisic acid (ABA) is a well-known compound in seed dormancy to which the embryo growth is very sensitive (Kermode 1995). Other than the above treatments, exogenous growth regulator treatments (i.e., gibberellins and cytokinins) have been found effective to break dormancy in many seed species (Koyuncu 2005; Bradford *et al.* 2008; Rita *et al.* 2010).

The studies of Hunt (2005) show that the seeds of *Heliotropium europaeum* are not dormant, while our primary study did not confirm his results. So, we checked the tests again with other seed lots and the results did not change. Therefore, these experiments are designed to overcome the possible seed dormancy of this plant.

MATERIALS AND METHODS

The seeds of *Heliotropium europaeum* used in this study are found throughout Maragheh and Bonab and occupy a range of semi-dryland habitats in East Azerbaijan province. Seeds were collected at maturity during the 2012 growing season. All seeds were air dried on the lab table at room temperature for approximately two weeks prior to the treatments. They were sorted by hand and air flowed to remove empty seeds in order to have a higher proportion of viable seeds in the experiments. Seed lot was separated to seven subsamples. One sample of seeds was stored at dry conditions during 12 months and was used for after-ripening tests. However, the other subsamples were immediately subjected to following seed dormancy breaking treatments: gibberellic acid, hot water, KNO₃, cold stratification and washing. Untreated seeds were used as the control.

The hot water treatment was performed through soaking the seeds in hot distilled water at 100°C for one min. After the required soaking period, the seeds were removed and cooled. In cold stratification, seeds were placed between rolled filter paper moistened with distilled water and stored in a refrigerator at $2 \pm 1^\circ\text{C}$ for 30 days. In the gibberellic acid treatments, seeds were immersed in solutions of GA 250 mg L⁻¹ for 12 hours. With KNO₃ treatment, seeds were soaked in 1.5 percent KNO₃ solution for one hour. During the washing treatment control seeds were soaked in tap water for 30min then put under running tap water for 30 min. After treatments seeds surface were dried and considered for the germination and seedling test. Seeds were placed between moistened rolled papers and incubated at $22 \pm 2^\circ\text{C}$ in the light. Both control and treated seeds were counted as germinated when the radicles were $\geq 2\text{mm}$. Another test was carried out to evaluate seedling properties using the same protocol for germination test but the test finished after 14 days. At the end of seedling test, radicles and shoots were cut from the cotyledons and then length of radicles and shoots were measured and mean seedling lengths were determined. The dry weights of seeding properties were negligible, so these data are not mentioned in the means Table 1. The vigor index was calculated by following equation: Vigor Index = (Germination percentage \times Seedling length).

Statistical Analysis

The statistical analyses of data, analyzed by SAS Ver. 9.1 software, included ANOVA and Duncan's multiple range test ($p < 0.05$).

RESULTS AND DISCUSSION

There was a significant effect of treatments on seed germination percentage $p < 0.05$. The treatments had different effects on seed germination (Table 1 and Figure 1). In comparison with control, washing and hot water treatments did not improve this trait. However, after-ripening, Gibberellic acid, KNO₃ and cold stratification remarkably released seed dormancy. Among after-ripening, Gibberellic acid and KNO₃ treatments differences were not significant, while cold stratification had significant differences with them. The highest germination percentage (93%) was recorded with cold stratification followed by KNO₃, after-ripening and Gibberellic acid, respectively.

Table 1. Means comparison of seed treatments effect on dormancy and seedling traits of *Heliotropium europaeum*.

	Germination percentage	Radicle length (mm)	Shoot length (mm)	Shoot to radicle ratio	Seedling length (mm)	Vigor index
After ripening	37 b	2.11 a	2.05 b	1.02 b	4.22 b	1.58 b
Control	0 c	-----	-----	-----	-----	-----
Gibberellic acid	35 b	1.48 b	3.20 a	2.36 ab	4.69 ab	1.68 b
Hot water	0 c	-----	-----	-----	-----	-----
KNO₃	41 b	1.57 ab	3.69 a	2.71 a	5.14 a	2.16 b
Stratification	93 a	1.11 b	2.44 b	2.23 ab	3.58 c	3.35 a
Washing	0 c	-----	-----	-----	-----	-----

Means with the same letter are not significantly different at $p < 0.05$.

The results of seedlings derived from effective treatments revealed that radicals, shoot, and seedling lengths varied among treatments (Table 1). Seedling raised in after-ripening treatment had longer radicals than those of the other treatments. However, this treatment produced shortest shoot lengths among treatments. Therefore the shoot to root ratio decreased (Table 1). Among treatments, potassium nitrate had superior shoot length. The highest seedling length was also recorded for KNO₃ treatments. The calculation of vigor index showed that the cold stratification had the highest index among treatments while it was not statistically significant among other treatments (Table 1).

After a period of 12 months storage, a significant increase in seed germination was observed and after-ripening increased seed germination to 37%. It seems that fresh seeds gradually release from dormancy during storage time or the intensity of their dormancy decreases among the seed lots. This plant has indeterminate growth and, as long as it lives, continues to add new flowers. Consequently, seeds with different ages are formed on an inflorescence. Seed age has shown to be effective in overcoming seed dormancy. Oziegbe *et al.* (2010) reported a high germination percentage in six months old seeds of *Ludwigia adscendens* L. as compared to freshly shed seeds. *Triticum durum* L. (Grilli 1993), *Bromus tectorum* L (Christensen *et al.* 1996; Meyer 2004) and *Avena fatua* (Foley, 1994) caryopses have a relative dormancy, which is released during a period of after-ripening. Dry after-ripening probably changes growth inhibitors content or growth promoter contents in the seed embryo, inducing the seed to germinate (Baskin and Baskin 2001). It has been suggested that during dry after-ripening, gene expression (β -1,3-glucanase transcription and translation) in *Nicotiana tabacum* L. (Solanaceae) seeds may be altered (Leubner-Metzger 2005). GA₃ broke the seed dormancy and 35% of seeds began to germinate. Gibberellins have been shown to break dormancy in numerous seed types (Bewley and Black 1994; Debeaujon *et al.* 2007). Debeaujon and Koornneef (2000) showed that dormancy and germination are the consequence of a balance between many stimulating and inhibiting factors, including GA and ABA. Exogenous application of GA₃ can alter the balance in favor of the stimulus. So, germination processes have been probably started with a high level of GA₃. The results show that seeds of common heliotrope germinate with 1.5% KNO₃ treatment. This finding is in agreement with Bian *et al.* (2013) findings which showed seed germination of *Sorbus pohuashanensis* improved with KNO₃ treatment. Numerous inorganic compounds are present in the soil matrix, however, only nitrate and nitrite significantly influence dormancy state (Baskin and Baskin 2001). Although, the application of KNO₃ is a simple treatment in many seed-testing laboratories, however a clear explanation for its action has not been studied.




Figure 1. Effects of treatments on seed germination of *Heliotrope*

Hot water did not overcome seed dormancy in *Heliotrope*. Though, this treatment has been reported to be effective for removing seed dormancy of wild *Vigna* species (Wang *et al.* 2011) and *Gymnocladus assamicus* (Choudhury *et al.* 2009). The ability of hot water to release seed germination appears to be depending on species, treatment duration and the availability and requirement of light after treatment. The experiment shows that the water content of *Heliotrope* seeds raise within first hour after immersing in water therefore the physical (seed coats hardness) dormancy cannot be a limiting factor for this plant (water content of seeds raised up to 50% during 30 min.). Hot water treatment is commonly used to release the seed coat imposing dormancy. Repeated washing of seeds did not improve their germination. In contrast to Hunt (2005) findings in *Heliotrope*, however, no evidence of leachable germination inhibitors was detected. Probably in our study, the given time for washing was not sufficient to remove the leachable germination inhibitors. The cold stratification vigorously broke dormancy status of *Heliotrope* and germination percentage increased to 93%. The findings of the current study are consistent with those of Koyuncu (2005) and Pipinis *et al.* (2011). These findings further support the idea of Baskin and Baskin (2001) which suggested dispersed seeds in autumn, commonly require cold stratification to stimulate germination. Dormancy would have been ecologically important for common *heliotrope*. It would inhibit germination of fresh seeds in late summer and early autumn when environmental conditions are unfavorable for successive plant growth and development.

CONCLUSIONS

We found that seeds of *Heliotropium europae* L. had a high level of dormancy at the fresh harvesting stage. It would be ecologically beneficial for summer annual weeds in North West of Iran, because it prevents fresh seeds from germinating at the end of the growing season when the low or freezing temperature would kill establishing seedlings during winter. Seed treatment with dry after- ripening, Gibberellic acid, KNO₃ and cold stratification can be effective in releasing dormancy and the most effective method to break seed dormancy was cold stratification.

REFERENCES

- Baskin C, and Baskin J (2001). Seeds Ecology, Biogeography and Evolution of Dormancy and Germination. Academic Press, London, UK.
- Bewley JD, Black M(1994). Seeds; Physiology of Development and Germination. Plenum Press, New York, USA.
- Bian L, Yang L, Wang J, and Shen H (2013). Effects of KNO₃ pretreatment and temperature on seed germination of *Sorbus pohuashanensis*. Journal of Forestry Research. 24:309-316.
- Bradford KJ, Benech-Arnold RL, Come D, and Corbineau F (2008). Quantifying the sensitivity of barley seed germination to oxygen, abscisic acid, and gibberellin using a population-based threshold model. Journal of Experimental Botany. 59: 335–347.
- Choudhury B, Khan BI, and Das AK (2009). Seed dormancy and germination in *Gymnocladus assamica*: An endemic legume tree from Northeast India. Seed Sci. and Technol. 37: 582-588.
- Christensen M, Meyer SE, and Allen PS (1996). A hydrothermal time model of seed after-ripening in *Bromus tectorum* L. Seed Science Research. 6: 155-163.
- Debeaujon I, and Koornneef M (2000). Gibberellin requirement for Arabidopsis thaliana seed germination is determined both by testa characteristics and embryonic ABA. Plant Physiol. 26:415–424.
- Debeaujon I, Lepiniec L, Pourcel L, and Routaboul JM (2007). Seed coat development and dormancy. In: Bradford K., Nonogaki H., editors. Seed development, dormancy and germination. 1. Vol. 6. Oxford, UK: Blackwell Publishing. pp. 25–43.
- Finch-Savage WE, and Leubner-Metzger G (2006). Seed dormancy and the control of germination. New Phytol. 171:501-23.
- Foley ME (1994). Temperature and water status of seed affect after-ripening in wild oat (*Avena fatua* L.). Weed Science. 42: 200-204.
- Francisco H, Dübbern DS, and Júlio MF (2001). The seed coat as a modulator of seed-environment relationships in Fabaceae. Rev. Bras. Bot. 24: 365- 375.
- Grilli I, Anguillesi MC, and Floris C (1993). Breaking of dormancy during after-ripening of *Triticum durum* DESF.: RNA, poly(A)+RNA and protein metabolism in imbibed embryos. Phytol. 34:119-130.
- Hunt JR (2005). The ecology of common heliotrope (*Heliotropium europaeum* L.) in a Mediterranean dry-land cropping system. PhD thesis, Faculty of Land and Food Resources, The University of Melbourne.
- Kermode AR (1995). Regulatory mechanisms in the transition from seed development to germination: interactions between the embryos and the seed environment. In: Kigel, J.; Galili, G. (Ed.) Seed development and germination. New York: Marcel Dekker, p.273-332.
- Kim SY, De Datta SK, and Mercado BL (1990). The effect of chemical and heat treatments on germination of *Commelina benghalensis* L. aerial seeds. Weed Res. 30:109–116.
- Koyuncu F (2005). Breaking seed dormancy in black mulberry (*Morus nigra* L.) By cold stratification and exogenous application of gibberellic acid. Acta Biologica Cracoviensia series Botanica. 47: 23–26.
- Leubner-Metzger G (2005). Beta-1,3-Glucanase gene expression in low-hydrated seeds as a mechanism for dormancy release during tobacco after-ripening. The Plant Journal. 41: 133-145.
- Meyer SE, Nelson DL, and Carlson SL (2004). Ecological genetics of vernalization response in *bromus tectorum* L. (Poaceae). Annals of Botany. 93: 653-663.
- Mohammadi G, Jalali S, Shirkhani A, and Shabani G (2012). Effects of seedhardness breaking techniques on Okra (*Abelmoschus esculentus* L.) germination. Intl J Agri Crop Sci. 4: 264-273.
- Oziegbe M, Faluyi JSO, and Oluwaranti A (2010). Effect of seed age and soil texture on the germination of some Ludwigia species (Onagraceae) in Nigeria. Acta Bot. Croat. 69: 249–257.
- Pipinis E, Milios E, Smiris P, and Gioumousidis C (2011). Effect of acid scarification and cold moist stratification on the germination of *Cercis siliquastrum* L. seeds. Turk J Agric For. 35: 259-264.
- Rita NK, Chakraborty S, and Kumar NJI (2010). Methods to break seed dormancy of *Andrographis paniculata* (Burm.f.Nees): An important medicinal herb of tropical asia. Advances In Bioresearch. 1: 35 – 39.
- Saeedi M, and Semnani M (2009). Chemical composition and antimicrobial activity of the essential oil of *Heliotropium europaeum*. Chemistry of Natural Compounds. 45:1, p98.
- Tung LD, and Serrano EP (2011). Effects of warm water in breaking dormancy of rice seed. Omonrice. 18: 129-136.
- Wang YR, Hanson J, Mariam YW (2011). Breaking hard seed dormancy in diverse accessions of five wild Vigna species by hot water and mechanical scarification. Seed Science and Technology. 39: 12-20.
- Watanabe H, Kasagaya Y, and Saigusa M (2002). Environmental factors affecting germination of apple of Peru. Weed Sci. 50:152–156.