

RESEARCH ARTICLE

J Res Vet Med. 2022; 41 (1) 43-48
DOI:10.30782/jrv.1059876

Immunohistochemical Investigation of TNF- α Expression in Sheep and Goat Lung Paraffin Blocks Infected with Natural Respiratory Syncytial Virus

• Funda Terzi^{1*}, • Yavuz Ulusoy², • Bahadır Kılınc²,
• Ayşe Gül Dal Tabag³, • Rabia Salik⁴

1 Kastamonu University, Faculty of Veterinary Medicine, Department of Pathology, 37150, Kastamonu, Turkey.

2 Etlik Central Veterinary Control and Research Institute, Pathology Department, 06020, Ankara, Turkey

3 Samsun Veterinary Control Institute, Pathology Department, 55200, Samsun, Turkey.

4 Selcuk University, Faculty of Veterinary Medicine, Department of Pathology,

Received 19-01-2022 Accepted 16-05-2022

Abstract

Respiratory Syncytial Virus (RSV) is an important viral agent of sheep and goat respiratory tract infections. RSV usually replicates in airway epithelium, inducing proinflammatory cytokines and chemokines. In this study, it was aimed to investigate TNF- α expression in natural RSV-infected sheep and goat lung paraffin blocks by immunohistochemical method. The material of the study consisted of 29 lung paraffin blocks (19 sheep and 10 goats), which were admitted to Etlik Veterinary Control and Research Institute with the suspicion of pneumonia from Ankara and surrounding provinces between 2015 and 2020. Histopathological findings such as degeneration and desquamation in the bronchial and bronchial epithelium, fibromuscular hypertrophy, hyperplasia in the peribronchial lymphoid tissue, cell infiltration in the interalveolar septum were common in sheep and goat lung paraffin block tissues. Immunohistochemically, RSV reaction was not statistically significant in sheep and goats in bronchial and bronchial epithelium and cell debris, bronchial glands, interalveolar septum inflammatory cells and cytoplasm of alveolar macrophages ($p > 0.05$). A statistically significant increase in TNF- α expression was determined in goat lung paraffin blocks compared to sheep ($p < 0.05$).

In conclusion, an increase in TNF α expression was detected in native RSV-infected sheep and goat lung tissue. It is thought that the results of the study will contribute to the treatment of sheep and goat respiratory tract RSV infections.

Keywords; RSV, TNF- α , Immunohistochemistry, Sheep, Goat

Introduction

Respiratory Syncytial Virus (RSV) targets the respiratory system such as rhinitis, bronchiolitis, pneumonia and sometimes causes otitis media¹. RSV is an enveloped, non-segmented, negative-stranded RNA virus in the family Paramyxoviridae²⁻⁴. RSVs have been defined as human RSV (HRSV), bovine RSV (BRSV), ovine RSV (ORSV), and caprine RSV (CRSV)⁵⁻⁷. RSV is the most common and important cause of lower respiratory tract infections

in cattle, calves, and young infants, and clinical and patho-logical findings in BRSV-infected calves and HRSV-infect-ed infants are similar^{8,9}. Sheep develop respiratory tract infections due to ORSV in the natural environment and are also susceptible to BRSV¹⁰. It has also been shown that BRSV, HRSV, and CRSV strains are closely related⁶. RSVs have three transmembrane glycoproteins of the 11 proteins encoded by the viral genome, and the large binding gly-coprotein (G), fusion protein (F), and small hydrophobic protein (SH) are found inside the viral envelope G (bind-

* Corresponding author: Funda Terzi, E-mail: fundaterzi@kastamonu.edu.tr, Kastamonu University, Faculty of Veterinary Medicine, Department of Pathology, Kuzeyskent Campus- TR- 37150, Kastamonu/ Türkiye, Tel: +9 0366 2805114 18, Fax: +9 0366 280 10 38

ing protein) and F (fusion protein) encoded by RSVs are the two major surface glycoproteins¹¹. Sheep RSV has the F gene and its encoded protein, bovine RSV has 85% nucleotide and 94% amino acid identity¹². The G and F proteins play important roles in the binding and entry of HRSV and BRSV into target cells¹³.

Transmission of RSV occurs through aerosol droplets of infected animals, direct contact, or indirectly through contaminated surfaces¹⁴. RSV proliferates in ciliary airway epithelial cells and type I and type II alveolar pneumocyte cells^{15,16} and shows necrotic bronchiolitis, degenerative or necrotic changes in bronchiolar epithelium, and peribronchial lymphocyte infiltration^{17,18}. Diffuse interstitial thickening of the alveolar septa is seen as a result of infiltration of inflammatory cells (ie, lymphocytes, macrophages) in the alveolar septum and type II pneumocyte hyperplasia. In the alveolar lumen, fluid with seroproteins, cell debris, a few alveolar macrophages, and occasionally neutrophils are detected¹⁹. Multinucleated syncytial cells are most often observed in the bronchiolar wall or lumen and less frequently in the alveoli⁵. Eosinophilic intracytoplasmic viral inclusions can be seen in mononuclear cells or epithelium as well as in syncytial cells²⁰.

Alveolar macrophages are a possible target for RSV infection²¹. In RSV infection, a rapid response develops by macrophages with the production of a wide variety of cytokines such as TNF- α , IL-1, IL-6, IL-8^{19,21,22}. Tumor necrosis factor-alpha (TNF- α) is a central mediator of immune and inflammatory responses and is important in host responses to infection²³. In addition, TNF- α against HRSV was determined to have an antiviral effect in vitro^{24,25}.

In this study, it was aimed to investigate TNF- α expression in lung paraffin blocks in sheep and goat natural RSV infections by immunohistochemical method.

Material and Method

The study material consisted of a total of 29 lung archive paraffin blocks, from 19 sheep and 10 goats, which were brought to xxx Institute with the suspicion of pneumonia between 2015 and 2020. In addition, histopathological findings of interstitial pneumonia and immunohistochemical RSV positive lung paraffin blocks were used. This study was approved by the Kastamonu University Animal Experiments Local Ethics Committee (Number: E-16498365-604.01.02-2100063448 Decision No: 25/3).

Histopathological Method

Lung paraffin blocks, 5 μ m thick sections with a microtome (Leica RM 2255) were taken on slides with poly-L-ly-

sine. Sections were stained on a Hematoxylin-eosin Stainer (Shandon) and covered with a coverslip by dripping entellan. Hematoxylin-Eosin stained sections were examined under a light microscope and photographed (Leica DM 400B). Histopathology scores were classified as mild (+), moderate (++) and severe (+++) expression according to the criteria determined by Grubor, et al.²⁶.

Immunohistochemical Method

Sections taken from lung paraffin blocks to adhesive slides, deparaffinization and rehydration procedures were applied. Immunohistochemical staining was performed according to the Mouse and Rabbit Specific HRP/DAB IHC Detection Kit-Micro polymer (ab236466) kit procedure. Hydrogen peroxide was dripped onto the sections. Proteinase K (ab64220) was added for antigen retrieval treatment and incubated at room temperature. The protein block solution was poured and left for 10 minutes and then incubated in primary antibody (1:100 anti Mouse RSV monoclonol (Cat NO: ab 43812), 1:350 anti- Rabbit TNF- α (Cat NO: ab 6671)) for 1 hour at room temperature. The Mouse Detection Reagent (Supplementary) solution was incubated for 10 min followed by a drop of Goat anti-rabbit HRP-conjugate for 15 min. DAB chromogen was added to demonstrate antigen-antibody reaction. Background staining was done with Mayer's hematoxylin. It was passed through xylol and alcohol series and closed with a coverslip by dripping enthalen. Sections were examined under a light microscope and photographed (Leica DM 400B). Immunohistochemical staining scores were classified as mild (+), moderate (++) and severe (+++) expression according to the criteria determined by Yavuz and Dinçel²⁷.

Statistics

Histopathological and immunohistochemical score in sheep and goat paraffin blocks were evaluated using IBM SPSS Statistics 25.0 software and Mann-Whitney's U test. The criterion for statistical significance was $p < 0.05$. All values are presented as mean \pm standard deviation.


Results

Histopathological Results

Histopathological findings of RSV positive cases in sheep and goat lung paraffin blocks were scored by semiquantitative method. Histopathological findings such as degeneration and desquamation in the bronchi and bronchiolar epithelium, fibromuscular hypertrophy (Fig. 1A), hyperplasia in the peribronchial lymphoid tissue (Fig. 1B), cell infiltration in the interalveolar septum (Fig. 1 C-D) were not statistically significant ($p > 0.05$) in sheep and goat lung paraffin block tissues (Table 1). It was determined that al-

veolar macrophages increased slightly in sheep and moderately increased in goats. In the interalveolar septum, syncytial giant cells were detected in 15 cases in sheep and 8 cases in goats.

Immunohistochemical Results




Figures 1. Hematoxylin-Eosin Staining. Lung tissue of sheep and goat. A. Moderate fibromuscular hypertrophy (black arrows) and bronchial gland hyperplasia. Bar:200 µm. B. Moderate peribronchiolar lymphoid tissue hyperplasia (black arrows), cell infiltration, and hyperemia in the interalveolar septum. Bar:200 µm. C. Moderate cell infiltration (black arrows) and hyperemia (red arrows) in the interalveolar septum. Bar:200 µm. D. Severe cell infiltration (black arrows) and hyperemia in the interalveolar septum. Bar: 50 µm.


In the Immunohistochemistry, RSV was found positive in lung paraffin block tissue of 19 sheep and 10 goats. Accordingly, RSV reaction in sheep and goat lungs, cell debris in bronchial and bronchiolar epithelium, and cell debris in lumen, bronchial glands and alveolar macrophages were statistically similar ($p>0.05$). While RSV antigen was found in the bronchial epithelium and cell debris in the lumen mild in 21% of and severe in 10% of sheep (Fig 2A), RSV antigen was not found in the bronchial epithelium and cell debris in the lumen of goats. RSV antigen was detected in bronchiol epithelium and cell debris in the lumen, mild (15% of sheep and 20% of goats) and severe only in 10% of sheep (Fig 2C). RSV antigen positive in bronchial glands was found only in one sheep (Fig 2B). Inter-alveolar septum cell infiltrations to RSV antigen in mild (31% of sheep and 50% of goats Fig 2D), moderate (21% of sheep, Fig 2E) and severe (15% of sheep and 20% of goats Fig 2F) was seen. In alveolar macrophages, RSV antigen was detected mild (57% of sheep, 60% of goats), moderate (15% and 10% of sheep), severe (15% of sheep and 30% of goats).

TNF-α expression in alveolar macrophages was mild (12

sheep and 3 goats Fig 3A), moderate (1 sheep and 3 goats Fig 3B), severe (2 sheep and 3 goats Fig 3C). It was determined that the expression of TNF-α in sheep and goat lung paraffin blocks was statistically more intensely stained ($p<0.05$) in goats than in sheep (Fig 4, Table 1).



Figures 2. Immunohistochemical staining of sheep and goat lung's. A. Severe anti-RSV reaction in the bronchial epithelium (black arrows). Bar: 100µm. B. Anti-RSV reaction (black arrows) in the bronchial glands. Bar: 100µm. C. Anti-RSV reaction in the bronchiolar epithelium (black arrows) and alveolar lumen. Bar: 100µm. D. Mild anti-RSV positive reaction (black arrows) in the cytoplasm of interalveolar septal cell infiltrates. Bar: 50µm. E. Moderate anti-RSV positive reaction (black arrows) in the interalveolar septum cell cytoplasm. Bar: 50µm. F. Severe anti-RSV positive reaction (black arrows) in the interalveolar septum cell cytoplasm. Bar: 50µm.



Figures 3. Immunohistochemical staining. Lungs. Sheep and goat. A. Mild anti-TNF-α positive reactions (black arrows) in the interalveolar septum cell cytoplasm. Bar:50µm. B. Moderate anti-TNF-α positive reaction (black arrows) in the interalveolar septum cell cytoplasm and alveolar macrophages. Bar: 50µm. C. Severe anti-TNF-α positive reaction (black arrows) in the interalveolar septum cell cytoplasm. Bar: 50µm.

Table 1. Statistical findings of histopathological and immunohistochemical staining scores in RSV antigen positive sheep and goat lung blocks.

	Lesions	Sheep	Goat	P value
Histopathology	Degeneration and desquamation of the bronchial and bronchiolar epithelium	1.21±0.91	1.50±0.84	0.438
	Fibromuscular hypertrophy	1.15±0.68	1.30±0.67	0.594
	Lymphoid hyperplasia	0.63±0.88	0.60±0.84	1.00
	Thickening of the interalveolar septum	1.89±0.65	1.50±0.53	0.122
	Epithelialization	1.00±0.33	1.20±0.63	0.242
	Hyalin membrane	0.84±0.37	1.10±0.56	0.162
	Intraalveolar edema	0.21±0.53	0.70±0.94	0.118
	Alveolar macrophage	1.89±0.93	2.10±0.56	0.496
	Hiperemi	1.42±0.60	1.20±0.42	0.334
	IHC	In the bronch epithelium and cell debris	0.52±0.96	0.00±0.00
In bronchial epithelium and cell debris		0.42±0.96	0.20±0.42	0.845
In the Bronchial Glands		0.36±0.95	0.00±0.00	0.193
In inflammatory cells in the interalveolar septal tissue.		1.10±0.99	1.10±1.10	0.903
Alveolar macrophages		1.36±0.89	1.70±0.94	0.379

Wan Witney U testi. Mean±Std. P<0.05.




Figure 4. TNF- α expression increased significantly in goat lung paraffin tissue compared to sheep ($p < 0.05$). Wan Witney U test. Mean+Std. * $p < 0.05$.

Discussion

Respiratory syncytial virus (RSV) is the leading cause of lower respiratory tract disease in humans and animals. RSV causes epidemics in ruminant herds and causes serious economic losses. RSV outbreaks usually occur in autumn and winter⁹. BRSV and HRSV proliferate in ciliary airway epithelial cells and type II pneumocytes^{15,17} and innate immunity is activated, resulting in the induction of proinflammatory cytokines and chemokines^{19,28,29}. In humans, calves and lambs, cytokines such as TNF- α , IL-6 and IL-8 increase gene expression from alveolar epithelium and alveolar macrophages³⁰⁻³². We investigated TNF- α expression in natural RSV-infected sheep and goat lung paraffin tissue by immunohistochemical method.

In RSV infections, cytokines and chemokines induce adhesion molecules^{19,28}, neutrophils and lymphocytes proliferation, resulting in bronchiolitis and interstitial pneumonia in the lung²⁹. Clinical symptoms such as fever, weakness, mucoid nasal discharge, continuous and deep breathing, tachypnea, cough, and anorexia are seen in RSV infections^{17,29,33,34}. Necropsy findings are similar in RSV infections of ruminants. Macroscopically, irregular lobular or fused atelectasis areas and consolidation areas localized in the cranioventral region of the lung are seen^{35,36}. In all lobes can be seen interstitial edema, while emphysema is detected in the diaphragm lobes³⁷. Also, the trachea, bronchi, and bronchioles may contain mucopurulent discharge^{5,17}. In RSV infections, histopathologically, bronchitis, bronchiolitis, thickening of inflammatory cell infiltrates in the alveolar septum, lymphocyte accumulations in the perivascular and peribronchiolar spaces, hyperplasia in the epithelial cells of the bronchi and bronchioles are caused³⁵. There are also neutrophils and macrophages in the lumen of the bronchi, bronchioles and alveoli and multinu-

cleated syncytial cells and hyaline membranes³². Similar to previous studies, in the paraffin tissue of the naturally RSV virus-infected sheep and goat lungs were detected degeneration and desquamation of the bronchial and bronchiolar epithelium, fibromuscular hypertrophy, lymphoid hyperplasia, cell infiltration in the interalveolar septum, epithelialization, and hyaline membrane such as interstitial pneumonia findings. Interstitial pneumonia findings in sheep and goat lung paraffin blocks were not statistically significant ($p > 0.05$) and RSV infections were similar in sheep and goats.

RSV replication occurs in both non-ciliary cells of the bronchiole and types II cells at the bronchiole/alveolar junction, where airflow and gas exchange is greatly impaired due to cell damage and inflammation in these areas³⁸. In the study, it was determined that RSV antigen in the bronchial epithelium was mild in 21% of sheep and severe in 10%, and RSV antigen was not found in the bronchial epithelium of goats. It was detected in the lung paraffin blocks of sheep and goats with mild severity in the bronchiolar epithelium (15% and 20%, respectively) and severe only in 10% of sheep. In both humans and lambs, RSV viral antigen was detected in bronchial and bronchiolar epithelial cells, as well as in type II pneumocytes^{15,18}. In this study, RSV antigen was detected in sheep and goat lung bronchial and bronchiolar epithelial cells, similar to RSV antigen determination studies in humans and lambs. According to Viuff, et al.¹⁶ found that although the number of mature alveolar macrophages increased during experimental BRSV infection in calves, they only rarely contained RSV antigen. In the study, generally mild RSV reaction was determined in alveolar macrophages in sheep and goats.

TNF- α has an important role in the pathogenesis of the disease³². TNF- α is a 17-kd, 157-amino acid cytokine secreted by many cells, including macrophages, monocytes, T cells, natural killer cells, and neutrophils³⁹. Franke, et al.⁴⁰ found that RSV-infected macrophages released TNF- α in a virus dose-dependent manner in a study inoculating murine alveolar macrophages with RSV. Røntved, et al.³² identified elevated TNF- α levels on days when severe lung lesions and clinical manifestations were exacerbated and BRSV-antigen levels were highest. In the study, TNF- α expression in alveolar macrophages was mild (12 sheep and 3 goats-), moderate (1 sheep and 3 goats-), severe (2 sheep and 3 goats). In addition, TNF- α expression in the cytoplasm of interalveolar septal cell infiltrates was found to be significantly higher ($p < 0.05$) in goats compared to sheep. These findings show that natural RSV infections of sheep and goats cause in the lung increased expression of TNF- α .

Conclusion

Viral infections cause the release of many proinflammatory cytokines in the lung tissue. In this study, an increase in the expression of the proinflammatory cytokine TNF- α was detected in the lung tissue of naturally RSV infected sheep and goats. In addition, the results of the study are thought to contribute to the treatment of RSV infections in ruminants.

Acknowledgements

This study was supported by the Kastamonu Scientific Research project (Project No: KÜ-BAP 01/2020-2).

References

- Sitthicharoenchai P, Alnajjar S, Ackermann MR. A model of respiratory syncytial virus (RSV) infection of infants in newborn lambs. *Cell and tissue research* 2020;380:313-324.
- Alansari H, Potgieter L. Molecular cloning and sequence analysis of the phosphoprotein, nucleocapsid protein, matrix protein and 22K (M2) protein of the ovine respiratory syncytial virus. *Journal of General Virology* 1994;75:3597-3601.
- Lehmkuhl H, Smith M, Cutlip R. Morphogenesis and structure of caprine respiratory syncytial virus. *Archives of virology* 1980;65:269-276.
- Kast JI, McFarlane AJ, Głobińska A, et al. Respiratory syncytial virus infection influences tight junction integrity. *Clinical & Experimental Immunology* 2017;190:351-359.
- Sacco RE, McGill J, Pillatzki AE, et al. Respiratory syncytial virus infection in cattle. *Veterinary pathology* 2014;51:427-436.
- Trudel M, Nadon F, Simard C, et al. Comparison of caprine, human and bovine strains of respiratory syncytial virus. *Archives of virology* 1989;107:141-149.
- Grubbs ST, Kania SA, Potgieter L. Prevalence of ovine and bovine respiratory syncytial virus infections in cattle determined with a synthetic peptide-based immunoassay. *Journal of veterinary diagnostic investigation* 2001;13:128-132.
- Baker JC. Human and bovine respiratory syncytial virus: immunopathologic mechanisms. *Veterinary Quarterly* 1991;13:47-59.
- Van der Poel W, Brand A, Kramps J, et al. Respiratory syncytial virus infections in human beings and in cattle. *Journal of Infection* 1994;29:215-228.
- Meyerholz DK, Grubor B, Fach SJ, et al. Reduced clearance of respiratory syncytial virus infection in a preterm lamb model. *Microbes and infection* 2004;6:1312-1319.
- Eleraky NZ, Kania SA, Evermann JF, et al. Comparison of targeting F and G protein genes to detect bovine and ovine respiratory syncytial viruses. *Journal of veterinary diagnostic investigation* 2003;15:277-280.
- Eleraky NZ, Kania SA, Potgieter LN. The ovine respiratory syncytial virus F gene sequence and its diagnostic application. *Journal of veterinary diagnostic investigation* 2001;13:455-461.
- Bem RA, Domachowske JB, Rosenberg HF. Animal models of human respiratory syncytial virus disease. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 2011;301:L148-L156.
- Hägglund S, Svensson C, Emanuelson U, et al. Dynamics of virus infections involved in the bovine respiratory disease complex in Swedish dairy herds. *The Veterinary Journal* 2006;172:320-328.
- Johnson JE, Gonzales RA, Olson SJ, et al. The histopathology of fatal untreated human respiratory syncytial virus infection. *Modern Pathology* 2007;20:108-119.
- Viuff B, Tjørnehøj K, Larsen LE, et al. Replication and clearance of respiratory syncytial virus: apoptosis is an important pathway of virus clearance after experimental infection with bovine respiratory syncytial virus. *The American journal of pathology* 2002;161:2195-2207.
- Viuff B, Uttenthal Å, Tegtmeier C, et al. Sites of replication of bovine respiratory syncytial virus in naturally infected calves as determined by in situ hybridization. *Veterinary Pathology* 1996;33:383-390.
- Larios Mora A, Detalle L, Van Geelen A, et al. Kinetics of respiratory syncytial virus (RSV) Memphis strain 37 (M37) infection in the respiratory tract of newborn lambs as an RSV infection model for human infants. *PLoS One* 2015;10:e0143580.
- Valarcher J-F, Taylor G. Bovine respiratory syncytial virus infection. *Veterinary research* 2007;38:153-180.
- Garg R, Latimer L, Wang Y, et al. Maternal immunization with respiratory syncytial virus fusion protein formulated with a novel combination adjuvant provides protection from RSV in newborn lambs. *Vaccine* 2016;34:261-269.
- Panuska JR, Cirino NM, Midulla F, et al. Productive infection of isolated human alveolar macrophages by respiratory syncytial virus. *The Journal of clinical investigation* 1990;86:113-119.
- Midulla F, Villani A, Panuska JR, et al. Respiratory syncytial virus lung infection in infants: immunoregulatory role of infected alveolar macrophages. *Journal of Infectious Diseases* 1993;168:1515-1519.
- Neuzil KM, Tang Y-W, Graham BS. Protective role of

- TNF- α in respiratory syncytial virus infection in vitro and in vivo. *The American journal of the medical sciences* 1996;311:201-204.
24. Cirino NM, Panuska JR, Villani A, et al. Restricted replication of respiratory syncytial virus in human alveolar macrophages. *Journal of general virology* 1993;74:1527-1537.
 25. Merolla R, Rebert NA, Tsiviste PT, et al. Respiratory syncytial virus replication in human lung epithelial cells: inhibition by tumor necrosis factor alpha and interferon beta. *American journal of respiratory and critical care medicine* 1995;152:1358-1366.
 26. Grubor B, Gallup JM, Meyerholz DK, et al. Enhanced surfactant protein and defensin mRNA levels and reduced viral replication during parainfluenza virus type 3 pneumonia in neonatal lambs. *Clinical and Vaccine Immunology* 2004;11:599-607.
 27. Yavuz O, Dinçel GÇ. Expression of COX-2, HMGB-1 and CD68 in lung tissue in sheep fibrinous bronchopneumonia. *Medycyna Weterynaryjna* 2020;76.
 28. F Rosenberg H, B Domachowske J. Inflammatory responses to respiratory syncytial virus (RSV) infection and the development of immunomodulatory pharmacotherapeutics. *Current medicinal chemistry* 2012;19:1424-1431.
 29. Guzman E, Taylor G. Immunology of bovine respiratory syncytial virus in calves. *Molecular immunology* 2015;66:48-56.
 30. Becker S, Quay J, Soukup J. Cytokine (tumor necrosis factor, IL-6, and IL-8) production by respiratory syncytial virus-infected human alveolar macrophages. *The Journal of Immunology* 1991;147:4307-4312.
 31. Sow FB, Gallup JM, Olivier A, et al. Respiratory syncytial virus is associated with an inflammatory response in lungs and architectural remodeling of lung-draining lymph nodes of newborn lambs. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 2011;300:L12-L24.
 32. Runtved C, Tjørnehøj K, Viuff B, et al. Increased pulmonary secretion of tumor necrosis factor- α in calves experimentally infected with bovine respiratory syncytial virus. *Veterinary Immunology and Immunopathology* 2000;76:214.
 33. Vilcek S, Elvander M, Ballagi-Pordany A, et al. Development of nested PCR assays for detection of bovine respiratory syncytial virus in clinical samples. *Journal of Clinical Microbiology* 1994;32:2225-2231.
 34. Gershwin LJ. Immunology of bovine respiratory syncytial virus infection of cattle. *Comparative immunology, microbiology and infectious diseases* 2012;35:253-257.
 35. Belknap EB, Ciszewski DK, Baker JC. Experimental respiratory syncytial virus infection in calves and lambs. *Journal of Veterinary Diagnostic Investigation* 1995;7:285-298.
 36. West K, Petrie L, Haines DM, et al. The effect of formalin-inactivated vaccine on respiratory disease associated with bovine respiratory syncytial virus infection in calves. *Vaccine* 1999;17:809-820.
 37. Kimman T, Straver P, Zimmer G. Pathogenesis of naturally acquired bovine respiratory syncytial virus infection in calves: morphologic and serologic findings. *American journal of veterinary research* 1989;50:684-693.
 38. Derscheid RJ, Ackermann MR. Perinatal lamb model of respiratory syncytial virus (RSV) infection. *Viruses* 2012;4:2359-2378.
 39. Krishnadasan B, Naidu BV, Byrne K, et al. The role of proinflammatory cytokines in lung ischemia-reperfusion injury. *The Journal of thoracic and cardiovascular surgery* 2003;125:261-272.
 40. Franke G, Freiherst J, Steinmüller C, et al. Interaction of alveolar macrophages and respiratory syncytial virus. *Journal of immunological methods* 1994;174:173-184.