Turkish Journal of Mathematics

Volume 38 | Number 2

Article 16

1-1-2014

On generalized Robertson--Walker spacetimes satisfying some curvature condition

KADRİ ARSLAN

RYSZARD DESZCZ

RIDVAN EZENTAŞ

MARIAN HOTLOS

CENGİZHAN MURATHAN

Follow this and additional works at: https://journals.tubitak.gov.tr/math

Part of the Mathematics Commons

Recommended Citation

ARSLAN, KADRİ; DESZCZ, RYSZARD; EZENTAŞ, RIDVAN; HOTLOS, MARIAN; and MURATHAN, CENGİZHAN (2014) "On generalized Robertson--Walker spacetimes satisfying some curvature condition," *Turkish Journal of Mathematics*: Vol. 38: No. 2, Article 16. https://doi.org/10.3906/mat-1304-3 Available at: https://journals.tubitak.gov.tr/math/vol38/iss2/16

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.



 ${\tt http://journals.tubitak.gov.tr/math/}$

Research Article

On generalized Robertson–Walker spacetimes satisfying some curvature condition

Kadri ARSLAN^{1,*}, Ryszard DESZCZ², Ridvan EZENTAȘ¹, Marian HOTLOŚ³, Cengizhan MURATHAN¹

¹Department of Mathematics, Art and Science Faculty, Uludağ University, Bursa, Turkey ²Department of Mathematics, Wrocław University, of Environmental, and Life Sciences 50-357 Wrocław, Poland ³Institute of Mathematics, and Computer Science, Wrocław University, of Technology, 50-350 Wrocław, Poland

Received: 02.04.2013	٠	Accepted: 19.10.2013	٠	Published Online: 27.01.2014	٠	Printed: 24.02.2014
----------------------	---	----------------------	---	------------------------------	---	----------------------------

Abstract: We give necessary and sufficient conditions for warped product manifolds (M, g), of dimension ≥ 4 , with 1-dimensional base, and in particular, for generalized Robertson–Walker spacetimes, to satisfy some generalized Einstein metric condition. Namely, the difference tensor $R \cdot C - C \cdot R$, formed from the curvature tensor R and the Weyl conformal curvature tensor C, is expressed by the Tachibana tensor Q(S, R) formed from the Ricci tensor S and R. We also construct suitable examples of such manifolds. They are quasi-Einstein, i.e. at every point of M rank $(S - \alpha g) \le 1$, for some $\alpha \in \mathbb{R}$, or non-quasi-Einstein.

Key words: Warped product, generalized Robertson–Walker spacetime, Einstein manifold, quasi-Einstein manifold, essentially conformally symmetric manifold, Tachibana tensor, generalized Einstein metric condition, pseudosymmetry type curvature condition, Ricci-pseudosymmetric hypersurface

1. Introduction

A semi-Riemannian manifold (M,g), $n = \dim M \ge 3$, is said to be an *Einstein manifold* if at every point its Ricci tensor S is proportional to the metric tensor g, i.e. on M we have

$$S = \frac{\kappa}{n} g, \qquad (1)$$

where κ is the scalar curvature of (M, g). In particular, if S vanishes identically on M then (M, g) is called a Ricci flat manifold. If at every point of M its Ricci tensor satisfies rank $S \leq 1$ then (M, g) is called a Ricci-simple manifold (see, e.g., [31, 41]).

Let (M,g), $n \ge 3$, be a semi-Riemannian manifold and let U_S be the set of all points of M at which $S \ne \frac{\kappa}{n}g$. The manifold (M,g), $n \ge 3$, is said to be quasi-Einstein (see, e.g., [56] and [22] and references therein) if at every point of $U_S \subset M$ we have rank $(S - \alpha g) = 1$, for some $\alpha \in \mathbb{R}$.

For the curvature tensor R and the Weyl conformal curvature tensor C of (M,g), $n \ge 4$, we can define on M the (0, 6)-tensors $R \cdot C$ and $C \cdot R$. For precise definition of the symbols used, we refer to Sections 2 and 3 of this paper, as well as to [5, 22, 24, 32]. It is obvious that for any Ricci flat, as well as conformally flat, semi-Riemannian manifold (M,g), $n \ge 4$, we have $R \cdot C - C \cdot R = 0$. For non-Ricci flat Einstein manifolds the

^{*}Correspondence: arslan@uludag.edu.tr

²⁰¹⁰ AMS Mathematics Subject Classification: Primary 53B20, 53B30, 53B50; Secondary 53C25, 53C50, 53C80.

tensor $R \cdot C - C \cdot R$ is nonzero. Namely, any Einstein manifold $(M, g), n \ge 4$, satisfies ([32], Theorem 3.1)

$$R \cdot C - C \cdot R = \frac{\kappa}{(n-1)n} Q(g,R) = \frac{\kappa}{(n-1)n} Q(g,C), \qquad (2)$$

i.e. at every point of M the difference tensor $R \cdot C - C \cdot R$ and the Tachibana tensor Q(g, R), or Q(g, C), are linearly dependent. We also mention that for any semi-Riemannian manifold (M, g), $n \ge 4$, we have some identity (see Eq. (32)) that expresses the tensor $R \cdot C - C \cdot R$ by some (0, 6)-tensors. In particular, by making use of that identity we can express the difference tensor of some hypersurfaces in space forms by a linear combination of the Tachibana tensors Q(g, R) and Q(S, R) ([24], Theorem 3.2; see also our Theorem 6.1(ii) and Proposition 4.1).

We can also investigate semi-Riemannian manifolds (M,g), $n \ge 4$, for which the difference tensor $R \cdot C - C \cdot R$ is expressed by one of the following Tachibana tensors: Q(g,R), Q(g,C), Q(S,R), or Q(S,C). In this way we obtain 4 curvature conditions. The first results related to those conditions are given in [32]. We refer to [22] for a survey on this subject. Since these conditions are satisfied on any semi-Riemannian Einstein manifold, they can be named generalized Einstein metric conditions (cf. [6], Chapter 16). In particular, (2) is also a condition of this kind and in Section 2 we present results on manifolds satisfying

$$R \cdot C - C \cdot R = L_1 Q(g, R). \tag{3}$$

In this paper we restrict our investigations to non-Einstein and nonconformally flat semi-Riemannian manifolds $(M,g), n \ge 4$, satisfying on $U = \{x \in M : Q(S,R) \neq 0 \text{ at } x\}$ the condition

$$R \cdot C - C \cdot R = LQ(S,R), \qquad (4)$$

where L is some function on this set. We recall that at all points of a semi-Riemannian manifold (M, g), $n \ge 3$, at which its Ricci tensor S is nonzero and Q(S, R) = 0, we have ([9], Theorem 4.1)

$$R \cdot R = 0, \tag{5}$$

i.e. such a manifold is semisymmetric. We also recall that if

$$R \cdot S = 0 \tag{6}$$

holds on a semi-Riemannian manifold then it is called *Ricci-semisymmetric*. The condition (4), under some additional assumptions, was considered in [45] (see Theorem 2.2 of this paper).

Our main results are related to warped products $\overline{M} \times_F \widetilde{N}$ with 1-dimensional base manifold $(\overline{M}, \overline{g})$ satisfying (4). Evidently, generalized Robertson–Walker spacetimes are warped products of this kind. We investigate separately 2 cases where the fiber $(\widetilde{N}, \widetilde{g})$ is either non-Einstein or Einstein manifold. In the first case, in Section 4, we prove that the associated function L satisfies $L = \frac{1}{n-2}$ and we show that the warping function Fis a polynomial of the second degree. Moreover, $(\widetilde{N}, \widetilde{g})$ satisfies some curvature condition presented by (46) (see Theorems 4.1 and 4.2). Furthermore, in the second case (see Section 5), i.e. when $\overline{M} \times_F \widetilde{N}$ is a quasi-Einstein manifold, we show that also $L = \frac{1}{n-2}$ and F is a polynomial of the second degree (see Theorem 5.1). Based on these results, in Section 6 we give examples of warped products satisfying (4). In particular, we construct an example of a warped product manifold satisfying (4) having non-Einstein fiber realizing (46). Finally, we mention that recently hypersurfaces in space forms having the tensor $R \cdot C - C \cdot R$ expressed by some Tachibana tensors were investigated in [26].

2. Preliminaries

Throughout this paper all manifolds are assumed to be connected paracompact manifolds of class C^{∞} . Let (M, g) be an *n*-dimensional semi-Riemannian manifold and let ∇ be its Levi-Civita connection and $\mathfrak{X}(M)$ the Lie algebra of vector fields on M. We define on M the endomorphisms $X \wedge_A Y$ and $\mathcal{R}(X, Y)$ of $\mathfrak{X}(M)$ by

$$\begin{aligned} &(X \wedge_A Y)Z &= A(Y,Z)X - A(X,Z)Y, \\ &\mathcal{R}(X,Y)Z &= \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]}Z, \end{aligned}$$

respectively, where A is a symmetric (0,2)-tensor on M and $X, Y, Z \in \mathfrak{X}(M)$. The Ricci tensor S, the Ricci operator S, and the scalar curvature κ of (M,g) are defined by $S(X,Y) = tr\{Z \to \mathcal{R}(Z,X)Y\}$, g(SX,Y) = S(X,Y), and $\kappa = trS$, respectively. The endomorphism $\mathcal{C}(X,Y)$ is defined by

$$\mathcal{C}(X,Y)Z = \mathcal{R}(X,Y)Z - \frac{1}{n-2}(X \wedge_g SY + SX \wedge_g Y - \frac{\kappa}{n-1}X \wedge_g Y)Z,$$

assuming that $n \ge 3$. Now the (0, 4)-tensor G, the Riemann-Christoffel curvature tensor R, and the Weyl conformal curvature tensor C of (M, g) are defined by

$$G(X_1, X_2, X_3, X_4) = g((X_1 \wedge_g X_2) X_3, X_4),$$

$$R(X_1, X_2, X_3, X_4) = g(\mathcal{R}(X_1, X_2) X_3, X_4),$$

$$C(X_1, X_2, X_3, X_4) = g(\mathcal{C}(X_1, X_2) X_3, X_4),$$

respectively, where $X_1, X_2, \ldots \in \mathfrak{X}(M)$. Furthermore, we define the following sets: $U_R = \{x \in M : R \neq \frac{\kappa}{(n-1)n}G \text{ at } x\}, U_S = \{x \in M : S \neq \frac{\kappa}{n}g \text{ at } x\}, \text{ and } U_C = \{x \in M : C \neq 0 \text{ at } x\}.$ It is easy to see that $U_S \cup U_C = U_R$.

Let $\mathcal{B}(X,Y)$ be a skew-symmetric endomorphism of $\mathfrak{X}(M)$ and let B be a (0,4)-tensor associated with $\mathcal{B}(X,Y)$ by

$$B(X_1, X_2, X_3, X_4) = g(\mathcal{B}(X_1, X_2) X_3, X_4).$$
(7)

The tensor B is said to be a generalized curvature tensor if

$$B(X_1, X_2, X_3, X_4) + B(X_2, X_3, X_1, X_4) + B(X_3, X_1, X_2, X_4) = 0,$$

$$B(X_1, X_2, X_3, X_4) = B(X_3, X_4, X_1, X_2).$$

Let $\mathcal{B}(X,Y)$ be a skew-symmetric endomorphism of $\mathfrak{X}(M)$ and let B be the tensor defined by (7). We extend the endomorphism $\mathcal{B}(X,Y)$ to derivation $\mathcal{B}(X,Y)$. of the algebra of tensor fields on M, assuming that it commutes with contractions and $\mathcal{B}(X,Y) \cdot f = 0$, for any smooth function f on M. Now for a (0,k)-tensor field $T, k \ge 1$, we can define the (0, k+2)-tensor $B \cdot T$ by

$$(B \cdot T)(X_1, \dots, X_k; X, Y) = (\mathcal{B}(X, Y) \cdot T)(X_1, \dots, X_k)$$

= $-T(\mathcal{B}(X, Y)X_1, X_2, \dots, X_k) - \dots - T(X_1, \dots, X_{k-1}, \mathcal{B}(X, Y)X_k).$

In addition, if A is a symmetric (0, 2)-tensor then we define the (0, k+2)-tensor Q(A, T), named a Tachibana tensor ([26]), by

$$Q(A,T)(X_1,...,X_k;X,Y) = (X \wedge_A Y \cdot T)(X_1,...,X_k)$$

= $-T((X \wedge_A Y)X_1,X_2,...,X_k) - \cdots - T(X_1,...,X_{k-1},(X \wedge_A Y)X_k).$

In this manner we obtain the (0, 6)-tensors $B \cdot B$ and Q(A, B). Setting in the above formulas $\mathcal{B} = \mathcal{R}$ or $\mathcal{B} = \mathcal{C}$, T = R or T = C or T = S, A = g or A = S, we get the tensors $R \cdot R$, $R \cdot C$, $C \cdot R$, $R \cdot S$, Q(g, R), Q(S, R), Q(g, C), and Q(g, S).

Let B_{hijk} , T_{hijk} , and A_{ij} be the local components of generalized curvature tensors B and T and a symmetric (0,2)-tensor A on M, respectively, where $h, i, j, k, l, m, p, q \in \{1, 2, ..., n\}$. The local components $(B \cdot T)_{hijklm}$ and $Q(A, T)_{hijklm}$ of the tensors $B \cdot T$ and Q(A, T) are the following:

$$(B \cdot T)_{hijklm} = g^{pq}(T_{pijk}B_{qhlm} + T_{hpjk}B_{qilm} + T_{hipk}B_{qjlm} + T_{hijp}B_{qklm}),$$

$$Q(A,T)_{hijklm} = A_{hl}T_{mijk} + A_{il}T_{hmjk} + A_{jl}T_{himk} + A_{kl}T_{hijm}$$
$$- A_{hm}T_{lijk} - A_{im}T_{hljk} - A_{jm}T_{hilk} - A_{km}T_{hijl}.$$
(8)

For a symmetric (0, 2)-tensor E and a (0, k)-tensor T, $k \ge 2$, we define their Kulkarni–Nomizu product $E \wedge T$ by (see, e.g., [21])

$$(E \wedge T)(X_1, X_2, X_3, X_4; Y_3, \dots, Y_k)$$

= $E(X_1, X_4)T(X_2, X_3, Y_3, \dots, Y_k) + E(X_2, X_3)T(X_1, X_4, Y_3, \dots, Y_k)$
 $-E(X_1, X_3)T(X_2, X_4, Y_3, \dots, Y_k) - E(X_2, X_4)T(X_1, X_3, Y_3, \dots, Y_k).$

According to [26], the tensor $E \wedge T$ is called a *Kulkarni–Nomizu tensor*. Clearly, the tensors R, C, G, and $E \wedge F$, where E and F are symmetric (0, 2)-tensors, are generalized curvature tensors.

A semi-Riemannian manifold (M,g), $n \ge 3$, is said to be locally symmetric if $\nabla R = 0$ holds on M. It is obvious that the last condition leads immediately to the integrability condition (5). Manifolds satisfying (5) are called semisymmetric ([58, 59]). Riemannian semisymmetric manifolds were classified in [58] and [59]. Non-Riemannian semi-Riemannian manifolds (M,g), $n \ge 4$, with parallel Weyl tensor ($\nabla C = 0$), which are in addition nonlocally symmetric ($\nabla R \ne 0$) and nonconformally flat ($C \ne 0$) are called essentially conformally symmetric manifolds, e.c.s. manifolds, in short ([13]–[18]). E.c.s. manifolds are semisymmetric (see, e.g., [13, 14]). In Remark 6.1(v) we present more details related to those manifolds. Another important subclass of semisymmetric manifolds, investigated recently, form second-order symmetric manifolds [7, 55], i.e. semi-Riemannian manifolds satisfying $\nabla \nabla R = 0$. As a weaker condition than (5) there is

$$R \cdot R = L_R Q(g, R) \,, \tag{9}$$

which is considered on $U_R \subset M$, and hence L_R is a function uniquely determined on this set. On $M \setminus U_R$ we have $R \cdot R = Q(g, R) = 0$. We note that Q(g, R) = 0 at a point if and only if $R = \frac{\kappa}{(n-1)n}G$ at this point. A semi-Riemannian manifold (M, g), $n \ge 3$, is said to be pseudosymmetric if (9) holds on $U_R \subset M$ [5, 19, 28, 43].

A semi-Riemannian manifold (M,g), $n \ge 3$, is said to be *Ricci-symmetric* if $\nabla S = 0$ holds on M. It is obvious that the last condition leads immediately to the integrability condition (6). Manifolds satisfying (6) are called *Ricci-semisymmetric*. As a weaker condition than (6) there is

$$R \cdot S = L_S Q(g, S) \,, \tag{10}$$

which is considered on $U_S \subset M$, and hence L_S is a function uniquely determined on this set. On $M \setminus U_S$ we have $R \cdot S = Q(g, S) = 0$. We note that Q(g, S) = 0 at a point if and only if (1) holds at this point (cf. [9], Lemma 2.1 (i)). A semi-Riemannian manifold (M, g), $n \ge 3$, is said to be *Ricci-pseudosymmetric* if (10) holds on $U_S \subset M$ [5, 19, 28, 46].

Every locally symmetric, resp. semisymmetric and pseudosymmetric, manifold is Ricci-symmetric, resp. Ricci-semisymmetric and Ricci-pseudosymmetric. In all cases, the converse statements are not true. We refer to [5, 22, 28, 44] for a wider presentation of results related to these classes of manifolds.

A geometric interpretation of (9), resp. (10), is given in [43], resp. in [46]. Semi-Riemannian manifolds for which their curvature tensor R is expressed by a linear combination of the Kulkarni–Nomizu tensors $S \wedge S$, $g \wedge S$, and G are called Roter-type manifolds; see, e.g., [36] and references therein. Precisely, a semi-Riemannian manifold (M, g), $n \ge 4$, is said to be a *Roter-type manifold* if

$$R = \frac{\phi}{2} S \wedge S + \mu g \wedge S + \eta G \tag{11}$$

holds on the set U_1 of all points of $U_S \cap U_C \subset M$ at which rank $(S - \alpha g) \ge 2$, for every $\alpha \in \mathbb{R}$. It is easy to prove that the functions ϕ , μ , and η are uniquely determined on U_1 . Using (11) and suitable definitions we can verify that on U_1 the condition (9) is satisfied (e.g., see [36], Eqs. (7) and (8); [22], Theorem 6.7) with $L_R = \phi^{-1}((n-2)(\mu^2 - \phi\eta) - \mu)$, and that the difference tensor $R \cdot C - C \cdot R$ is expressed on U_1 by a linear combination of the tensors Q(S, R), Q(g, R), and Q(S, G) ([24], Eq. (47)), or, equivalently, by a linear combination of the tensors Q(g, R) and Q(S, G) ([24], Eq. (48)).

Semi-Riemannian manifolds (M,g), $n \ge 4$, satisfying (3) on $U_S \cap U_C \subset M$ were investigated in [31]. Among other results in that paper it was proven that: (i) $R \cdot C = C \cdot R = 0$ and rank S = 1 hold on $U_S \cap U_C$, provided that (M,g) is a quasi-Einstein manifold; and (ii) (11), with some special coefficients ϕ, μ, η such that $R \cdot R = 0$, and $C \cdot R = -L_1 Q(g, R)$ hold on $U_S \cap U_C$, provided that (M,g) is a non-quasi-Einstein manifold. We also mention that manifolds satisfying

$$C \cdot R = LQ(g,R) \tag{12}$$

were investigated in [50]. Furthermore, we have

Theorem 2.1 ([32], Theorem 4.1 and Corollary 4.1) Let (M,g), $n \ge 4$, be a semi-Riemannian manifold. If $R \cdot C - C \cdot R = LQ(g,C)$ holds on $U_S \cap U_C \subset M$, for some function L, then $R \cdot R = LQ(g,R)$ and $C \cdot R = 0$ on this set. In particular, if $R \cdot C = C \cdot R$ holds on $U_S \cap U_C$ then $R \cdot R = R \cdot C = C \cdot R = 0$ on this set.

As we remarked in Section 1, manifolds satisfying (4) were investigated in [45]. We have

Theorem 2.2 ([45], Theorem 3.1 and Proposition 3.1) (i) Let (M, g), $n \ge 4$, be a nonconformally flat and non-Einstein Ricci-semisymmetric manifold satisfying (4). Then on the set consisting of all points of Mat which L is nonzero we have $L = \frac{1}{n-2}$ and

$$R(\mathcal{S}X, Y, Z, W) = \frac{\kappa}{n-1} R(X, Y, Z, W).$$
(13)

(ii) Let (M,g), $n \ge 4$, be a semi-Riemannian manifold satisfying (13). Then (M,g) is a Ricci-semisymmetric manifold fulfilling (4) with $L = \frac{1}{n-2}$.

We mention that hypersurfaces isometrically immersed in spaces of constant curvature having the tensor $R \cdot C$ expressed by a linear combination of the Tachibana tensors Q(g,R), Q(S,R), and Q(S,G) were investigated in [40].

Quasi-Einstein hypersurfaces isometrically immersed in spaces of constant curvature were investigated in [33, 41]; see also the references therein. In particular, in [33] an example of a quasi-Einstein hypersurface in a semi-Riemannian space of constant curvature was found. More precisely, in that paper it was shown that some warped product $\overline{M} \times_F \widetilde{N}$, with dim $\overline{M} = 1$ and dim $\widetilde{N} \ge 4$, can be locally realized as a nonpseudosymmetric Ricci-pseudosymmetric quasi-Einstein hypersurface in a semi-Riemannian space of constant curvature. The difference tensor of that hypersurface is expressed by a linear combination of the tensors Q(g, R) and Q(S, R).

3. Warped product manifolds

Warped products play an important role in Riemannian geometry (see, e.g., [49, 51]) as well as in general relativity theory (see, e.g., [3, 4, 34, 51]). Many well-known spacetimes of this theory, i.e. solutions of the Einstein field equations, are warped products, e.g., the Schwarzschild, Kottler, Reissner–Nordström, Reissner–Nordström–de Sitter, and Vaidya, as well as Robertson–Walker, spacetimes. We recall that a warped product $\overline{M} \times_F \widetilde{N}$ of a 1-dimensional manifold $(\overline{M}, \overline{g})$, $\overline{g}_{11} = -1$, and a 3-dimensional Riemannian space of constant curvature $(\widetilde{N}, \widetilde{g})$, with a warping function F, is said to be a Robertson–Walker spacetime (see, e.g., [3, 4, 51, 57]). It is well-known that the Robertson–Walker spacetimes are conformally flat quasi-Einstein manifolds. More generally, one also considers warped products $\overline{M} \times_F \widetilde{N}$ of $(\overline{M}, \overline{g})$, $dim \overline{M} = 1$, $\overline{g}_{11} = -1$, with a warping function F and (n-1)-dimensional Riemannian manifold $(\widetilde{N}, \widetilde{g})$, $n \ge 4$. Such warped products are called generalized Robertson–Walker spacetimes [1, 2, 38, 53, 54]. Curvature conditions of pseudosymmetry type on such spacetimes have been considered among others in [8, 9, 10, 12, 35, 36, 37, 42, 48]. We also mention that Einstein generalized Robertson–Walker spacetimes were classified in [2]. From (1) and (2) we immediately get the following:

Theorem 3.1 On any Einstein manifold (M,g), $n \ge 4$ we have

$$R \cdot C - C \cdot R = \frac{1}{n-1} Q(S,R).$$
(14)

In particular, (14) holds on any Einstein generalized Robertson–Walker spacetime.

Let now $(\overline{M}, \overline{g})$ and $(\widetilde{N}, \widetilde{g})$, dim $\overline{M} = p$, dim $\widetilde{N} = n - p$, $1 \leq p < n$, be semi-Riemannian manifolds. Let $F : \overline{M} \to \mathbb{R}^+$ be a positive smooth function on \overline{M} . The warped product manifold, or in short warped product, $\overline{M} \times_F \widetilde{N}$ of $(\overline{M}, \overline{g})$ and $(\widetilde{N}, \widetilde{g})$ is the product manifold $\overline{M} \times \widetilde{N}$ with the metric $g = \overline{g} \times_F \widetilde{g}$ defined by $\overline{g} \times_F \widetilde{g} = \pi_1^* \overline{g} + (F \circ \pi_1) \pi_2^* \widetilde{g}$, where $\pi_1 : \overline{M} \times \widetilde{N} \longrightarrow \overline{M}$ and $\pi_2 : \overline{M} \times \widetilde{N} \longrightarrow \widetilde{N}$ are the natural projections on \overline{M} and \widetilde{N} , respectively. With respect to Corollary 3.1, in this paper we consider non-Einstein warped products $\overline{M} \times_F \widetilde{N}$ with 1-dimensional base manifold $(\overline{M}, \overline{g})$ and an (n-1)-dimensional fiber $(\widetilde{N}, \widetilde{g}), n \ge 4$.

Let $\{\overline{U} \times \widetilde{V}; x^1, x^2 = y^1, \ldots, x^n = y^{n-1}\}$ be a product chart for $\overline{M} \times \widetilde{N}$, where $\{\overline{U}; x^1\}$ and $\{\widetilde{V}; y^\alpha\}$ are systems of charts on $(\overline{M}, \overline{g})$ and $(\widetilde{N}, \widetilde{g})$, respectively. The local components of the metric $g = \overline{g} \times_F \widetilde{g}$ with respect to this chart are the following: $g_{11} = \overline{g}_{11} = \varepsilon = \pm 1$, $g_{hk} = F \widetilde{g}_{\alpha\beta}$ if $h = \alpha$ and $k = \beta$, and $g_{hk} = 0$ otherwise, $\alpha, \beta, \gamma, \cdots \in \{2, \ldots, n\}$ and $h, i, j, k \ldots \in \{1, 2, \ldots, n\}$. We will denote by bars (resp., by tildes) tensors formed from \overline{g} (resp., \widetilde{g}). It is known that the local components Γ_{ij}^h of the Levi-Civita connection ∇ of $\overline{M} \times_F \widetilde{N}$ are the following (see, e.g., [49, 35]):

$$\Gamma_{11}^{1} = 0, \quad \Gamma_{\beta\gamma}^{\alpha} = \widetilde{\Gamma}_{\beta\gamma}^{\alpha}, \quad \Gamma_{\alpha\beta}^{1} = -\frac{\varepsilon}{2} F' \widetilde{g}_{\alpha\beta},$$

$$\Gamma_{\alpha\beta}^{\alpha} = \frac{1}{2F} F' \delta_{\beta}^{\alpha}, \quad \Gamma_{\alpha1}^{1} = \Gamma_{11}^{\alpha} = 0, \quad F' = \partial_{1}F = \frac{\partial F}{\partial x^{1}}.$$
 (15)

The local components R_{hijk} of the curvature tensor R and the local components S_{hk} of the Ricci tensor S of $\overline{M} \times_F \widetilde{N}$, which may not vanish identically, are the following (see, e.g., [20, 35]):

$$R_{\alpha 11\beta} = -\frac{1}{2} T_{11} \tilde{g}_{\alpha\beta} = -\frac{\operatorname{tr} T}{2} g_{11} \tilde{g}_{\alpha\beta} , \quad R_{\alpha\beta\gamma\delta} = F(\tilde{R}_{\alpha\beta\gamma\delta} - \frac{\Delta_1 F}{4F} \tilde{G}_{\alpha\beta\gamma\delta}) , \quad (16)$$

$$S_{11} = -\frac{n-1}{2F} T_{11}, \quad S_{\alpha\beta} = \tilde{S}_{\alpha\beta} - \left(\frac{\operatorname{tr} T}{2} + (n-2)\frac{\Delta_1 F}{4F}\right) \tilde{g}_{\alpha\beta}, \tag{17}$$

$$T_{11} = F'' - \frac{(F')^2}{2F}, \quad \text{tr } T = \bar{g}^{11}T_{11} = \varepsilon \left(F'' - \frac{(F')^2}{2F}\right),$$

$$\Delta_1 F = \Delta_{1\bar{g}}F = \bar{g}^{11}(F')^2 = \varepsilon (F')^2. \tag{18}$$

The scalar curvature κ of $\overline{M} \times_F \widetilde{N}$ satisfies the following relation:

$$\kappa = \frac{1}{F} \Big(\tilde{\kappa} - (n-1) \Big(\operatorname{tr} T + (n-2) \frac{\Delta_1 F}{4F} \Big) \Big).$$
⁽¹⁹⁾

Using (8), (16), and (17) we can check that the local components $Q(g, R)_{hijklm}$ and $Q(S, R)_{hijklm}$ of the Tachibana tensors Q(g, R) and Q(S, R), which may not vanish identically, are the following:

$$Q(g,R)_{1\beta\gamma\delta1\mu} = Fg_{11}\left(\widetilde{R}_{\mu\beta\gamma\delta} + \left(\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F}\right)\widetilde{G}_{\mu\beta\gamma\delta}\right),\tag{20}$$

$$Q(g,R)_{\alpha\beta\gamma\delta\lambda\mu} = F^2 Q(\tilde{g},\tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}, \qquad (21)$$

ARSLAN et al./Turk J Math

$$Q(S,R)_{1\beta\gamma\delta1\mu} = -\frac{\operatorname{tr} T}{2}g_{11}\left((n-1)\widetilde{R}_{\mu\beta\gamma\delta} - \widetilde{g}_{\beta\gamma}\widetilde{S}_{\delta\mu} + \widetilde{g}_{\beta\delta}\widetilde{S}_{\gamma\mu} + \left(\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F}\right)\widetilde{G}_{\mu\beta\gamma\delta}\right), \qquad (22)$$

$$Q(S,R)_{1\beta 1\delta\lambda\mu} = -\frac{\operatorname{tr} T}{2} g_{11} Q(\tilde{g}, \tilde{S})_{\beta\delta\lambda\mu}, \qquad (23)$$

$$Q(S,R)_{\alpha\beta\gamma\delta\lambda\mu} = FQ(\widetilde{S},\widetilde{R})_{\alpha\beta\gamma\delta\lambda\mu} - \frac{\Delta_1 F}{4}Q(\widetilde{S},\widetilde{G})_{\alpha\beta\gamma\delta\lambda\mu} - F\left(\frac{\operatorname{tr} T}{2} + \frac{(n-2)\Delta_1 F}{4F}\right)Q(\widetilde{g},\widetilde{R})_{\alpha\beta\gamma\delta\lambda\mu}.$$
(24)

Let V be the (0,4)-tensor with the local components

$$V_{hijk} = g^{lm} S_{hl} R_{mijk} = S_h^{\ l} R_{lijk} \,. \tag{25}$$

Using (16) and (17) we can verify that the only nonzero components of this tensor are the following:

$$V_{1\beta\gamma1} = \frac{n-1}{4F} (\operatorname{tr} T) T_{11} \tilde{g}_{\beta\gamma} = \frac{n-1}{4F} (\operatorname{tr} T)^2 g_{11} \tilde{g}_{\beta\gamma},$$

$$V_{\alpha11\delta} = -\frac{\operatorname{tr} T}{2F} g_{11} \Big(\widetilde{S}_{\alpha\delta} - \Big(\frac{\operatorname{tr} T}{2} + \frac{(n-2)\Delta_1 F}{4F} \Big) \tilde{g}_{\alpha\delta} \Big),$$

$$V_{\alpha\beta\gamma\delta} = \widetilde{S}_{\alpha}^{\ \epsilon} \widetilde{R}_{\epsilon\beta\gamma\delta} - \Big(\frac{\operatorname{tr} T}{2} + \frac{(n-2)\Delta_1 F}{4F} \Big) \widetilde{R}_{\alpha\beta\gamma\delta} - \frac{\Delta_1 F}{4F} \Big(\widetilde{g}_{\beta\gamma} \widetilde{S}_{\alpha\delta} - \widetilde{g}_{\beta\delta} \widetilde{S}_{\alpha\gamma} \Big)$$

$$+ \Big(\frac{\operatorname{tr} T}{2} + \frac{(n-2)\Delta_1 F}{4F} \Big) \frac{\Delta_1 F}{4F} \widetilde{G}_{\alpha\beta\gamma\delta}.$$
(26)

The last equality yields

$$V_{\alpha\beta\gamma\delta} + V_{\beta\alpha\gamma\delta} = \widetilde{S}_{\alpha}^{\ \epsilon} \widetilde{R}_{\epsilon\beta\gamma\delta} + \widetilde{S}_{\beta}^{\ \epsilon} \widetilde{R}_{\epsilon\alpha\gamma\delta} - \frac{\Delta_1 F}{4F} (\widetilde{g}_{\beta\gamma} \widetilde{S}_{\alpha\delta} - \widetilde{g}_{\beta\delta} \widetilde{S}_{\alpha\gamma} + \widetilde{g}_{\alpha\gamma} \widetilde{S}_{\beta\delta} - \widetilde{g}_{\alpha\delta} \widetilde{S}_{\beta\gamma}) = (\widetilde{R} \cdot \widetilde{S})_{\alpha\beta\gamma\delta} - \frac{\Delta_1 F}{4F} Q(\widetilde{g}, \widetilde{S})_{\alpha\beta\gamma\delta}.$$

$$(27)$$

Let P be a (0, 6)-tensor with local components

$$P_{hijklm} = g_{hl}V_{mijk} - g_{hm}V_{lijk} - g_{il}V_{mhjk} + g_{im}V_{lhjk} + g_{jl}V_{mkhi} - g_{jm}V_{lkhi} - g_{kl}V_{mjhi} + g_{km}V_{ljhi} - g_{ij}(V_{hklm} + V_{khlm}) - g_{hk}(V_{ijlm} + V_{jilm}) + g_{ik}(V_{hjlm} + V_{jhlm}) + g_{hj}(V_{iklm} + V_{kilm}).$$
(28)

The local components of P, which may not vanish identically, are the following:

$$P_{1\beta 1\delta\lambda\mu} = g_{11} \left((\widetilde{R} \cdot \widetilde{S})_{\beta\delta\lambda\mu} + (\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F}) Q(\widetilde{g}, \widetilde{S})_{\beta\delta\lambda\mu} \right),$$

$$P_{1\beta\gamma\delta1\mu} = g_{11} \left(\widetilde{S}_{\mu}^{\ \epsilon} \widetilde{R}_{\epsilon\beta\gamma\delta} - (\frac{\operatorname{tr} T}{2} + \frac{(n-2)\Delta_1 F}{4F}) \widetilde{R}_{\mu\beta\gamma\delta} + (\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F}) (\widetilde{g}_{\beta\gamma} \widetilde{S}_{\delta\mu} - \widetilde{g}_{\beta\delta} \widetilde{S}_{\gamma\mu}) + ((n-2)(\frac{\Delta_1 F}{4F})^2 - \frac{(\operatorname{tr} T)^2}{4} - \frac{(n-3)\operatorname{tr} T}{2} \frac{\Delta_1 F}{4F}) \widetilde{G}_{\mu\beta\gamma\delta} \right),$$

$$(29)$$

$$P_{\alpha\beta\gamma\delta\lambda\mu} = F\left(\tilde{g}_{\alpha\lambda}V_{\mu\beta\gamma\delta} - \tilde{g}_{\alpha\mu}V_{\lambda\beta\gamma\delta} - \tilde{g}_{\beta\lambda}V_{\mu\alpha\gamma\delta} + \tilde{g}_{\beta\mu}V_{\lambda\alpha\gamma\delta} + \tilde{g}_{\gamma\lambda}V_{\mu\delta\alpha\beta} - \tilde{g}_{\gamma\mu}V_{\lambda\delta\alpha\beta} - \tilde{g}_{\delta\lambda}V_{\mu\gamma\alpha\beta} + \tilde{g}_{\delta\mu}V_{\lambda\gamma\alpha\beta} - \tilde{g}_{\beta\gamma}(V_{\alpha\delta\lambda\mu} + V_{\delta\alpha\lambda\mu}) - \tilde{g}_{\alpha\delta}(V_{\beta\gamma\lambda\mu} + V_{\gamma\beta\lambda\mu}) + \tilde{g}_{\beta\delta}(V_{\alpha\gamma\lambda\mu} + V_{\gamma\alpha\lambda\mu}) + \tilde{g}_{\alpha\gamma}(V_{\beta\delta\lambda\mu} + V_{\delta\beta\lambda\mu})\right).$$
(31)

4. Warped products with non-Einsteinian fiber

Since we investigate non-Einstein and nonconformally flat manifolds satisfying (4), we restrict our considerations to the set $\mathcal{U} = U \cap U_S \cap U_C$.

We assume that the warped product $\overline{M} \times_F \widetilde{N}$ satisfies (4) and the fiber $(\widetilde{N}, \widetilde{g})$ is not Einsteinian. Now we shall use the following identity, which holds on any semi-Riemannian manifold ([24], Section 4):

$$(n-2)(R \cdot C - C \cdot R)_{hijklm} = Q(S,R)_{hijklm} - \frac{\kappa}{n-1}Q(g,R)_{hijklm} + P_{hijklm}.$$
(32)

Thus, in view of (32) and the definition of the tensor P, condition (4) can be written in the following form:

$$\left((n-2)L-1\right)Q(S,R)_{hijklm} = P_{hijklm} - \frac{\kappa}{n-1}Q(g,R)_{hijklm}.$$
(33)

For h = 1, $i = \beta$, j = 1, $k = \delta$, $l = \lambda$, $m = \mu$, in view of (20)–(24), (33) yields

$$\left((n-2)L-1\right)Q(S,R)_{1\beta 1\delta\lambda\mu} = P_{1\beta 1\delta\lambda\mu}.$$
(34)

Substituting (23) and (29) into (34) we obtain

$$(\widetilde{R} \cdot \widetilde{S})_{\beta\delta\lambda\mu} = \left(\frac{\Delta_1 F}{4F} - \frac{(n-2)L}{2} \operatorname{tr} T\right) Q(\widetilde{g}, \widetilde{S})_{\beta\delta\lambda\mu} \,. \tag{35}$$

On the other hand, (4) implies

$$(R \cdot C - C \cdot R)(X_1, X_2, X_3, X_4; X, Y) + (R \cdot C - C \cdot R)(X, Y, X_1, X_2; X_3, X_4) + (R \cdot C - C \cdot R)(X_3, X_4, X, Y; X_1, X_2) = 0,$$

which in virtue of Proposition 4.1 of [24] is equivalent to

$$(R \cdot C)(X_1, X_2, X_3, X_4; X, Y) + (R \cdot C)(X, Y, X_1, X_2; X_3, X_4)$$

+ $(R \cdot C)(X_3, X_4, X, Y; X_1, X_2) = 0.$

Furthermore, we have ([8], Section 3, eq. (3.19)):

$$(\widetilde{R} \cdot \widetilde{S})_{\beta\delta\lambda\mu} = \left(\frac{\Delta_1 F}{4F} - \frac{\operatorname{tr} T}{2}\right) Q(\widetilde{g}, \widetilde{S})_{\beta\delta\lambda\mu} \,. \tag{36}$$

Since (\tilde{N}, \tilde{g}) , dim $\tilde{N} \ge 3$, is not Einsteinian, the tensor $Q(\tilde{g}, \tilde{S})$ is a nonzero tensor. Let $Q(\tilde{g}, \tilde{S}) \ne 0$ at $x \in \mathcal{U}$. Thus, on a coordinate neighborhood $V_1 \subset \mathcal{U}$ of x, in virtue of (35) and (36), we get

$$\left(L - \frac{1}{n-2}\right) \operatorname{tr} T = 0.$$

We assert that tr T = 0. Supposing that tr $T \neq 0$ at $y \in V_1$ we have $L = \frac{1}{n-2}$ on some neighborhood $U_1 \subset V_1$ of y. Therefore, (33) reduces on U_1 to

$$P = \frac{\kappa}{n-1} Q(g,R) \,. \tag{37}$$

Evidently, on U_1 we also have

$$\frac{\Delta_1 F}{4F} - \frac{\operatorname{tr} T}{2} = const. \tag{38}$$

Now (37) gives

$$P_{1\beta\gamma\delta1\mu} = \frac{\kappa}{n-1} Q(g,R)_{1\beta\gamma\delta1\mu} \,. \tag{39}$$

Substituting into this equality (19), (22), and (30) we obtain

$$\widetilde{S}_{\mu}^{\ \epsilon} \widetilde{R}_{\epsilon\beta\gamma\delta} = \left(\frac{\widetilde{\kappa}}{n-1} - \frac{\operatorname{tr} T}{2}\right) \widetilde{R}_{\mu\beta\gamma\delta} + \left(\frac{\Delta_1 F}{4F} - \frac{\operatorname{tr} T}{2}\right) \left(\widetilde{g}_{\beta\gamma} \widetilde{S}_{\mu\delta} - \widetilde{g}_{\beta\delta} \widetilde{S}_{\mu\gamma}\right) - \left(\frac{\widetilde{\kappa}}{n-1} - \frac{\operatorname{tr} T}{2}\right) \left(\frac{\Delta_1 F}{4F} - \frac{\operatorname{tr} T}{2}\right) \widetilde{G}_{\mu\beta\gamma\delta} .$$

$$(40)$$

Using now (38) and (40) we see that tr T = const. and consequently also $\frac{\Delta_1 F}{4F} = const.$ on U_1 . Whence, after standard calculations, we deduce that F must be of the form

$$F(x^1) = (ax^1 + b)^2, \ a, b \in \mathbb{R}.$$
 (41)

For such F we have tr T = 0 on U_1 , a contradiction. Therefore

$$\operatorname{tr} T = 0 \tag{42}$$

on V_1 . Thus, (38) reduces on V_1 to

$$\frac{\Delta_1 F}{4F} = const. = c_1. \tag{43}$$

Note that (42) and (43), in the same manner as above, imply (41) and we have

$$\operatorname{tr} T = 0, \quad \frac{\Delta_1 F}{4F} = c_1 = \varepsilon a^2.$$
(44)

We prove now that

$$L = \frac{1}{n-2} \tag{45}$$

on V_1 . Applying (19), (20), (21), (30), (42), and (43) to

$$\left((n-2)L-1\right)Q(S,R)_{1\beta\gamma\delta1\mu} = P_{1\beta\gamma\delta1\mu} - \frac{\kappa}{n-1}Q(g,R)_{1\beta\gamma\delta1\mu}$$

we get

$$\widetilde{S}_{\mu}^{\ \epsilon}\widetilde{R}_{\epsilon\beta\gamma\delta} = \frac{\widetilde{\kappa}}{n-1}\,\widetilde{R}_{\mu\beta\gamma\delta} + \varepsilon a^2(\widetilde{g}_{\beta\gamma}\widetilde{S}_{\mu\delta} - \widetilde{g}_{\beta\delta}\widetilde{S}_{\mu\gamma}) - \frac{\varepsilon\widetilde{\kappa}a^2}{n-1}\,\widetilde{G}_{\mu\beta\gamma\delta}\,. \tag{46}$$

On the other hand, (26), by (42) and (43), gives

$$V_{\alpha\beta\gamma\delta} = \widetilde{S}_{\alpha}^{\ \epsilon} \widetilde{R}_{\epsilon\beta\gamma\delta} - (n-2)c_1\widetilde{R}_{\alpha\beta\gamma\delta} - c_1(\widetilde{g}_{\beta\gamma}\widetilde{S}_{\alpha\delta} - \widetilde{g}_{\beta\delta}\widetilde{S}_{\alpha\gamma}) + (n-2)c_1^2\widetilde{G}_{\alpha\beta\gamma\delta} \,,$$

which by (46) turns into

$$V_{\alpha\beta\gamma\delta} = \left(\frac{\tilde{\kappa}}{n-1} - (n-2)c_1\right)\tilde{R}_{\alpha\beta\gamma\delta} + (n-2)c_1^2\tilde{G}_{\alpha\beta\gamma\delta}.$$

Substituting this into (31) we obtain

$$P_{\alpha\beta\gamma\delta\lambda\mu} = F\left(\frac{\tilde{\kappa}}{n-1} - (n-2)c_1\right) Q(\tilde{g},\tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} \,. \tag{47}$$

Now

$$((n-2)L-1)Q(S,R)_{\alpha\beta\gamma\delta\lambda\mu} = P_{\alpha\beta\gamma\delta\lambda\mu} - \frac{\kappa}{n-1}Q(g,R)_{\alpha\beta\gamma\delta\lambda\mu},$$

by making use of (21), (47), and

$$\frac{\kappa}{n-1} = \frac{1}{F} \left(\frac{\tilde{\kappa}}{n-1} - (n-2)c_1 \right),$$

turns into

$$((n-2)L-1)Q(S,R)_{\alpha\beta\gamma\delta\lambda\mu} = 0.$$
(48)

Since $V_1 \subset \mathcal{U}$ and in virtue of (22), (23), and (42) we have

$$Q(S,R)_{1\beta\gamma\delta1\mu} = Q(S,R)_{1\beta1\delta\lambda\mu} = 0,$$

at least one of the local components of $Q(S, R)_{\alpha\beta\gamma\delta\lambda\mu}$ must be nonzero. Therefore, (48) implies (45). Thus we have proven:

Theorem 4.1 Let $\overline{M} \times_F \widetilde{N}$ be a warped product manifold with 1-dimensional base manifold $(\overline{M}, \overline{g})$ and non-Einstein (n-1)-dimensional fiber $(\widetilde{N}, \widetilde{g})$, $n \ge 4$. If (4) is satisfied on $\overline{M} \times_F \widetilde{N}$ then on the set \mathcal{U} we have (46) and

$$L = \frac{1}{n-2}, \quad F(x^1) = (ax^1 + b)^2, \ a, b \in \mathbb{R}.$$
(49)

Corollary 4.1 Let $\overline{M} \times_F \widetilde{N}$ be a generalized Robertson–Walker spacetime with non-Einstein fiber $(\widetilde{N}, \widetilde{g})$, $n \ge 4$. If (4) is satisfied on $\overline{M} \times_F \widetilde{N}$ then (46) and (49) hold on \mathcal{U} .

Proposition 4.1 Under assumptions of Theorem 4.1 the fiber manifold (\tilde{N}, \tilde{g}) is a Ricci-pseudosymmetric manifold of constant type (see, e.g., [41]), precisely

$$\widetilde{R} \cdot \widetilde{S} = \varepsilon a^2 Q(\widetilde{g}, \widetilde{S}) \,. \tag{50}$$

Moreover, if $n \ge 5$ then the difference tensor $\widetilde{R} \cdot \widetilde{C} - \widetilde{C} \cdot \widetilde{R}$ of the fiber is expressed by the Tachibana tensors $Q(\widetilde{S}, \widetilde{R})$ and $Q(\widetilde{g}, \widetilde{R})$; precisely, we have

$$(n-3)(\widetilde{R}\cdot\widetilde{C}-\widetilde{C}\cdot\widetilde{R}) = Q(\widetilde{S},\widetilde{R}) - \frac{\widetilde{\kappa}}{(n-1)(n-2)}Q(\widetilde{g},\widetilde{R}).$$
(51)

Proof First we observe that (46) implies (50). Applying the identity (32) to (\tilde{N}, \tilde{g}) we have

$$(n-3)(\widetilde{R}\cdot\widetilde{C}-\widetilde{C}\cdot\widetilde{R}) = Q(\widetilde{S},\widetilde{R}) + \widetilde{P} - \frac{\widetilde{\kappa}}{n-2}Q(\widetilde{g},\widetilde{R}).$$
(52)

Using now (46) we get

$$\widetilde{V}_{lphaeta\gamma\delta} + \widetilde{V}_{etalpha\gamma\delta} = arepsilon a^2 Q(\widetilde{g},\widetilde{S})_{lphaeta\gamma\delta}$$

and

$$\widetilde{P} = \frac{\widetilde{\kappa}}{n-1} Q(\widetilde{g}, \widetilde{R}) - \varepsilon a^2 Q(\widetilde{S}, \widetilde{G}) - \varepsilon a^2 \, \widetilde{g} \wedge Q(\widetilde{g}, \widetilde{S}) \,,$$

which by making use of $\tilde{g} \wedge Q(\tilde{g}, \tilde{S}) = -Q(\tilde{S}, \tilde{G})$ (see (28) of [24]) reduces to $\tilde{P} = \frac{\tilde{\kappa}}{n-1} Q(\tilde{g}, \tilde{R})$. Substituting this equality into (52) we obtain (51).

We have also the converse statement to Theorem 4.1.

Theorem 4.2 Let $(\overline{M}, \overline{g})$, $\overline{g}_{11} = \varepsilon$, be a 1-dimensional manifold and let $(\widetilde{N}, \widetilde{g})$ be an (n-1)-dimensional non-Einstein manifold, $n \ge 4$, satisfying (46). If $F(x^1) = (ax^1 + b)^2$, then the warped product $\overline{M} \times_F \widetilde{N}$ fulfills (4) with $L = \frac{1}{n-2}$.

Proof As we have seen (cf. (33)), (4) for $L = \frac{1}{n-2}$ takes the form

$$P = \frac{\kappa}{n-1} Q(g, R) \,. \tag{53}$$

Using now (27), (44), and (50) we have

$$V_{\alpha\beta\gamma\delta} + V_{\beta\alpha\gamma\delta} = 0.$$
⁽⁵⁴⁾

Taking into account (26), (44), and (46), we obtain

$$V_{\alpha\beta\gamma\delta} = \phi \widetilde{R}_{\alpha\beta\gamma\delta} + \psi \, \widetilde{G}_{\alpha\beta\gamma\delta}$$

for some ψ and $\phi = \frac{\tilde{\kappa}}{n-1} - \varepsilon(n-2)a^2$. Substituting the above equality and (54) into (31) we get

$$P_{\alpha\beta\gamma\delta\lambda\mu} = F\phi Q(\tilde{g},\tilde{R})_{\alpha\beta\gamma\delta\lambda\mu},$$

which in view of (19) and (44) takes the form

$$P_{\alpha\beta\gamma\delta\lambda\mu} = \frac{\kappa}{n-1} Q(g,R)_{\alpha\beta\gamma\delta\lambda\mu} \,.$$

Using (29), (44), and (50), we have $P_{1\beta 1\delta\lambda\mu} = 0$, which means that (34) is satisfied. Finally, in the same manner we obtain (39). Thus we see that (53) is satisfied for all systems of indices.

Corollary 4.2 The equality (46) is satisfied on every Einstein manifold (\tilde{N}, \tilde{g}) . Thus every warped product $\overline{M} \times_F \tilde{N}$ with 1-dimensional base (\overline{M}, \bar{g}) , Einsteinian fiber (\tilde{N}, \tilde{g}) , dim $\tilde{N} \ge 4$, which is not a space of constant curvature, and the warping function $F(x^1) = (ax^1 + b)^2$, $\varepsilon a^2 \neq \frac{\tilde{\kappa}}{(n-2)(n-1)}$, satisfies (4) with $L = \frac{1}{n-2}$.

Remark 4.1. Let $\overline{M} \times_F \widetilde{N}$ be the warped product manifold with 1-dimensional base $(\overline{M}, \overline{g})$, Einsteinian fiber $(\widetilde{N}, \widetilde{g})$, dim $\widetilde{N} \ge 4$, and the warping function $F(x^1) = (ax^1 + b)^2$, $\varepsilon a^2 = \frac{\widetilde{\kappa}}{(n-2)(n-1)}$. Now (17) yields S = 0, i.e. $\overline{M} \times_F \widetilde{N}$ is a Ricci-flat manifold.

5. Warped products with Einsteinian fiber

In this section we consider non-Einstein warped products $\overline{M} \times_F \widetilde{N}$, dim $\overline{M} = 1$, assuming that a fiber $(\widetilde{N}, \widetilde{g})$ is an Einstein manifold, i.e.

$$\widetilde{S}_{\alpha\beta} = \frac{\widetilde{\kappa}}{n-1} \, \widetilde{g}_{\alpha\beta} \,. \tag{55}$$

Using (17) we can easily show that such warped product is a quasi-Einstein manifold. It is worth noticing that $\widetilde{R} \neq \frac{\widetilde{\kappa}}{(n-1)(n-2)}\widetilde{G}$ on \mathcal{U} . Using (23), (29), and (55), we get

$$Q(S,R)_{1\beta 1\delta\lambda\mu} = P_{1\beta 1\delta\lambda\mu} = 0.$$
(56)

Analogously, in view of (22), (30), and (55), we have

$$Q(S,R)_{1\alpha\beta\gamma1\delta} = \frac{\operatorname{tr} T}{2} g_{11} \left(-(n-1)\widetilde{R}_{\delta\alpha\beta\gamma} + \left(\frac{\widetilde{\kappa}}{n-1} - \left(\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F}\right)\right) \widetilde{R}_{\delta\alpha\beta\gamma} \right),$$
(57)

$$P_{1\beta\gamma\delta1\mu} = g_{11} \left(\eta \widetilde{R}_{\mu\beta\gamma\delta} \right)$$
(58)

$$+ \left(\left(\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F}\right) \frac{\tilde{\kappa}}{n-1} + (n-2) \left(\frac{\Delta_1 F}{4F}\right)^2 - \frac{(\operatorname{tr} T)^2}{4} - (n-3) \frac{\operatorname{tr} T}{2} \frac{\Delta_1 F}{4F} \right) \widetilde{G}_{\mu\beta\gamma\delta} \right)$$

where

$$\eta = \frac{\tilde{\kappa}}{n-1} - \frac{\operatorname{tr} T}{2} - \frac{(n-2)\Delta_1 F}{4F}.$$

Finally, making use of (26) and (55), we obtain

$$V_{\alpha\beta\gamma\delta} = \eta \left(\widetilde{R}_{\alpha\beta\gamma\delta} - \frac{\Delta_1 F}{4F} \, \widetilde{G}_{\alpha\beta\gamma\delta} \right),$$

and next, in virtue of (24) and (31), also

$$Q(S,R)_{\alpha\beta\gamma\delta\lambda\mu} = F\eta Q(\tilde{g},\tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}, \qquad (59)$$

$$P_{\alpha\beta\gamma\delta\lambda\mu} = F\eta Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} \,. \tag{60}$$

Thus, taking into account (20), (21), and (56)-(60), we see that (33) is equivalent to the following 2 equalities:

$$\left((n-2)L-1\right)\frac{\operatorname{tr} T}{2}\left(-(n-1)\widetilde{R}+\left(\frac{\widetilde{\kappa}}{n-1}-\left(\frac{\operatorname{tr} T}{2}-\frac{\Delta_1 F}{4F}\right)\right)\widetilde{G}\right)$$
$$=\frac{\operatorname{tr} T}{2}\left(\widetilde{R}+\left(\frac{\operatorname{tr} T}{2}-\frac{\Delta_1 F}{4F}\right)\widetilde{G}\right),$$
(61)

$$\left((n-2)L-1\right)\eta Q(\tilde{g},\tilde{R}) = \frac{\operatorname{tr} T}{2} Q(\tilde{g},\tilde{R}).$$
(62)

We consider 2 cases: (i) tr T = 0 and (ii) tr $T \neq 0$.

(i) tr T = 0. Since $Q(\tilde{g}, \tilde{R}) \neq 0$ on \mathcal{U} , (62) leads to

$$\left((n-2)L-1\right)\left(\frac{\tilde{\kappa}}{n-1}-\frac{(n-2)\Delta_1 F}{4F}\right)=0.$$

Supposing that $\frac{\tilde{\kappa}}{(n-2)(n-1)} = \frac{\Delta_1 F}{4F}$ and using (17) we obtain S = 0, a contradiction. Thus we get $\frac{\tilde{\kappa}}{(n-2)(n-1)} \neq \frac{\Delta_1 F}{4F}$ and $L = \frac{1}{n-2}$. Moreover, solving the differential equation tr T = 0, one can see that the warping function F must be of the form (41). Thus, we have the situation described in Corollary 4.2.

(ii) tr $T \neq 0$. Now (61) leads to

$$(n-2)\big((n-1)L-1\big)\,\widetilde{R} = \Big((n-2)L\big(\frac{\widetilde{\kappa}}{n-1} - (\frac{\operatorname{tr} T}{2} - \frac{\Delta_1 F}{4F})\big) - \frac{\widetilde{\kappa}}{n-1}\Big)\widetilde{G}\,,$$

whence $L = \frac{1}{n-1}$ and

$$\frac{\Delta_1 F}{4F} - \frac{\operatorname{tr} T}{2} = \frac{\tilde{\kappa}}{(n-1)(n-2)}.$$
(63)

It is worth noticing that under the above equalities (62) also holds. Applying (63) to (17), after standard calculations, we get (1), a contradiction. Thus we have proven:

Theorem 5.1 Let $\overline{M} \times_F \widetilde{N}$ be a non-Einstein warped product manifold with 1-dimensional base manifold $(\overline{M}, \overline{g})$ and Einsteinian (n-1)-dimensional fiber $(\widetilde{N}, \widetilde{g})$. If (4) is satisfied on $\overline{M} \times_F \widetilde{N}$, then on the set \mathcal{U} we have:

$$L = \frac{1}{n-2}, \quad F(x^{1}) = (ax^{1}+b)^{2}, \ a, b \in \mathbb{R}, \quad \varepsilon a^{2} \neq \frac{\widetilde{\kappa}}{(n-2)(n-1)}$$

Remark 5.1. Let $\overline{M} \times_F \widetilde{N}$ be the warped product manifold with 1-dimensional base $(\overline{M}, \overline{g})$, Einsteinian fiber $(\widetilde{N}, \widetilde{g})$, dim $\widetilde{N} \ge 4$, and the warping function F.

(i) We assume that F satisfies (63). As we stated above, now $\overline{M} \times_F \widetilde{N}$ is an Einstein manifold. Furthermore, it is easy to check that (63), in view of (18), takes the form

$$FF'' - (F')^2 + 2\varepsilon C_1 F = 0, \quad C_1 = \frac{\tilde{\kappa}}{(n-1)(n-2)}.$$
 (64)

This is exactly equation (29) of [36]. We can check that the following functions are solutions of (64) (cf. [36], Lemma 3.1):

$$\begin{split} F(x^{1}) &= \varepsilon C_{1} \left(x^{1} + \frac{\varepsilon c}{C_{1}}\right)^{2}, \quad \varepsilon C_{1} > 0, \\ F(x^{1}) &= \frac{c}{2} \left(\exp(\pm \frac{b}{2}x^{1}) - \frac{2\varepsilon C_{1}}{b^{2}c} \exp(\mp \frac{b}{2}x^{1})\right)^{2}, \quad c > 0, \quad b \neq 0, \\ F(x^{1}) &= \frac{2\varepsilon C_{1}}{c} \left(1 + \sin(cx^{1} + b)\right), \quad \frac{\varepsilon C_{1}}{c} > 0, \end{split}$$

where b and c are constants and x^1 belongs to a suitable nonempty open interval of \mathbb{R} .

(ii) As was mentioned in Section 3, Einstein generalized Robertson–Walker spacetimes were classified in [2]. As was stated in [2], if $\overline{M} \times_f \widetilde{N}$ is a generalized Robertson–Walker spacetime, with $\varepsilon = -1$, the warping function f, and the Einsteinian fiber $(\widetilde{N}, \widetilde{g})$ and $\varepsilon = -1$ is an Einstein manifold, then the differential equations given in (3) of [2] must be satisfied. Those equations, adopted to our denotations, takes the form

$$ff'' = \frac{\kappa}{(n-1)n}f^2$$
, $\frac{\kappa}{(n-1)n}f^2 = \frac{\widetilde{\kappa}}{(n-2)(n-1)} + (f')^2$,

respectively. If now we set $f = \sqrt{F}$ then the equations presented above lead to (64).

6. Examples

Corollary 4.2 and Theorem 5.1 give rise to examples of warped products satisfying (4) with Einstein fiber. The problem of finding of a warped product satisfying (4) with non-Einstein fiber reduces, via Theorem 4.2, to the problem of finding an example of a semi-Riemannian manifold (\tilde{N}, \tilde{g}) , dim $\tilde{N} = n - 1 \ge 3$, fulfilling (46). To obtain a suitable example we will use results of [21, 24, 30]. First of all, we adopt results contained in Theorem 3.1 of [21] and in Theorem 3.2 of [24]. Those results we can present in the following:

Theorem 6.1 Let $(\widetilde{N}, \widetilde{g})$ be a hypersurface isometrically immersed in a semi-Riemannian space of constant curvature $N_s^n(c)$, $n \ge 4$, with signature (s, n - s), where $c = \frac{\tau}{(n-1)n}$, τ is the scalar curvature of the ambient space and \widetilde{g} is the metric tensor induced on \widetilde{N} . Moreover, let the second fundamental tensor H of \widetilde{N} satisfy on some nonempty connected set $\widetilde{U} \subset \widetilde{N}$ the equation

$$H^3 = tr(H) H^2 + \lambda H, \tag{65}$$

where λ is some function on \widetilde{U} , and let the constant $\varepsilon = \pm 1$ be defined by the Gauss equation of \widetilde{N} in $N_s^n(c)$, *i.e.* by

$$\widetilde{R} \; = \; \frac{\varepsilon}{2} \, H \wedge H + \frac{\tau}{(n-1)n} \, \widetilde{G} \, .$$

(i) (cf. [21], Theorem 3.1) On \widetilde{U} we have

$$\widetilde{S}_{\mu}^{\ \epsilon}\widetilde{R}_{\epsilon\beta\gamma\delta} = \mu\left(\widetilde{R}_{\mu\beta\gamma\delta} - \frac{\tau}{(n-1)n}\widetilde{G}_{\mu\beta\gamma\delta}\right) + \frac{\tau}{(n-1)n}\left(\widetilde{g}_{\beta\gamma}\widetilde{S}_{\mu\delta} - \widetilde{g}_{\beta\delta}\widetilde{S}_{\mu\gamma}\right),\tag{66}$$

where $\mu = \frac{(n-2)\tau}{(n-1)n} - \varepsilon \lambda$.

(ii) (cf. [24], Theorem 3.2) If $n \ge 5$ then on \widetilde{U} we have

$$(n-3)\left(\widetilde{R}\cdot\widetilde{C}-\widetilde{C}\cdot\widetilde{R}\right) = Q(\widetilde{S},\widetilde{R}) + \left(\frac{(n-2)\tau}{(n-1)n} - \varepsilon\,\lambda - \frac{\widetilde{\kappa}}{n-2}\right)Q(\widetilde{g},\widetilde{R}), \tag{67}$$

where $\tilde{\kappa}$ is the scalar curvature of \tilde{N} .

We note that (66) implies immediately that $\widetilde{R} \cdot \widetilde{S} = \frac{\tau}{(n-1)n} Q(\widetilde{g}, \widetilde{S})$. In addition, if we assume that on \widetilde{U} we have

$$\lambda = 0 \quad \text{and} \quad (n-2)\tau = n\widetilde{\kappa},$$
(68)

then (46) holds on \widetilde{U} . The last remark suggests a solution of our problem. Namely, the last 2 conditions are realized on the hypersurface presented in Example 5.1 of [30]. Let (M, g) be the manifold defined in Example 5.1 of [30]. We denote it by $(\widetilde{N}, \widetilde{g})$. Clearly $(\widetilde{N}, \widetilde{g})$ is a manifold of dimension ≥ 4 . However, it is easy to verify that if we repeat the construction of $(\widetilde{N}, \widetilde{g})$ for the 3-dimensional case then all curvature properties remain true, excluding, of course, properties expressed by its Weyl conformal curvature tensor. Thus, without loss of generality, we can assume that dim $\widetilde{N} = n - 1 \geq 3$. In Example 5.1 of [30], among other things, it was shown that $(\widetilde{N}, \widetilde{g})$ is locally isometric to a hypersurface in a semi-Riemannian space of nonzero constant curvature. Since our considerations are local, we can assume that $(\widetilde{N}, \widetilde{g})$ is a hypersurface isometrically immersed in that space. Since $(\widetilde{N}, \widetilde{g})$ fulfils (68), Theorem 4.2 finishes our construction. We note that by making use of (67) and (68), we obtain (51).

Remark 6.1. (i) The Roter-type warped products $\overline{M} \times_F \widetilde{N}$ with 1-dimensional base manifold $(\overline{M}, \overline{g})$ and non-Einstein (n-1)-dimensional fiber $(\widetilde{N}, \widetilde{g})$, $n \ge 4$, were investigated in [36]. Among other results it was proven that the curvature tensor \widetilde{R} of the fiber $(\widetilde{N}, \widetilde{g})$ is expressed by the Kulkarni–Nomizu tensors $\widetilde{S} \wedge \widetilde{S}$, $\widetilde{g} \wedge \widetilde{S}$, and \widetilde{G} , i.e. the fiber also is a Roter-type manifold, provided that $n \ge 5$. Therefore, if we assume that the fiber manifold $(\widetilde{N}, \widetilde{g})$ considered in Theorem 6.1 is a nonpseudosymmetric Ricci-pseudosymmetric hypersurface (for instance, the Cartan hypersurfaces of dimension 6, 12, or 24 have this property (see, e.g., [41])), then fibers of both constructions are nonisometric.

(ii) From (12), by a suitable contraction, we get

$$C \cdot S = LQ(g,S). \tag{69}$$

We refer to [47] and [50] for examples of warped products satisfying (69). The condition (69) holds on some hypersurfaces in semi-Riemannian space forms, and, in particular, on the Cartan hypersurfaces ([21], Theorems 3.1 and 4.3). Hypersurfaces in semi-Euclidean space satisfying (69) were investigated in [52]. (iii) We also can investigate semi-Riemannian manifolds (M, g), $n \ge 4$, satisfying on $U_C \subset M$ the following condition of pseudosymmetric type (see, e.g., [11, 29]):

$$R \cdot R - Q(S, R) = LQ(g, C), \qquad (70)$$

where L is some function on this set. Warped products satisfying (70) were investigated in [11]. Among other results, in [11] it was shown that this condition is satisfied on every 4-dimensional warped product $\overline{M} \times_F \widetilde{N}$ with 1-dimensional base. Thus, in particular, every 4-dimensional generalized Robertson–Walker spacetime satisfies (70). We mention that (70) holds on every hypersurface in a semi-Riemannian space of constant curvature (see, e.g., [22], eq. (4.4)).

(iv) In [23] (Example 4.1) a warped product $\overline{M} \times_F \widetilde{N}$ of an (n-1)-dimensional base $(\overline{M}, \overline{g}), n \ge 4$, and an 1-dimensional fiber $(\widetilde{N}, \widetilde{g})$ satisfying rank S = 1, $\kappa = 0$, $R \cdot R = 0$, and $C \cdot S = 0$ was constructed. In addition, we can easily check that (4) with $L = \frac{1}{n-2}$ and Q(S,C) = Q(S,R) hold on $\overline{M} \times_F \widetilde{N}$ ([25]). Therefore, on $\overline{M} \times_F \widetilde{N}$ we also have $(n-2)(R \cdot C - C \cdot R) = Q(S,C)$. Semi-Riemannian manifolds satisfying $R \cdot C - C \cdot R = LQ(S,C)$, for some function L, were investigated in [25]. An example of a quasi-Einstein non-Ricci-simple manifold satisfying the last condition was given in Section 6 of [22].

(v) We recall that non-Riemannian semi-Riemannian manifolds (M,g), $n \ge 4$, with parallel Weyl tensor $(\nabla C = 0)$, which are in addition nonlocally symmetric $(\nabla R \ne 0)$ and nonconformally flat $(C \ne 0)$, are called essentially conformally symmetric manifolds, or e.c.s. manifolds, in short (see, e.g., [13, 14]). E.c.s. manifolds are semisymmetric manifolds satisfying ([13], Theorems 7, 8 and 9): Q(S,C) = 0, $C(\mathcal{S}X,Y,Z,W) = 0$, $S(\mathcal{S}X,Y) = 0$, $\kappa = 0$. In addition, on every e.c.s. manifold (M,g) we have [14]: rank $S \le 2$ and $FC = (1/2) S \land S$, where F is some function on M, called the fundamental function. The local structure of e.c.s. manifolds is determined [15, 17]. Certain e.c.s. metrics are realized on compact manifolds [16, 18]. Let now (M,g), $n \ge 4$, be an e.c.s. manifold satisfying rank $S \le 1$. Now it is easy to check that at all points of M at which rank S = 1 the conditions Q(S,C) = 0, $C(\mathcal{S}X,Y,Z,W) = 0$ turn into Q(S,R) = 0, $R(\mathcal{S}X,Y,Z,W) = 0$, respectively. The last equality means that the tensor V, defined by (25), vanishes. Therefore, (28) reduces to P = 0. Now we see that the identity (32) turns into $R \cdot C - C \cdot R = 0$, and, in consequence, at all points of M at which rank S = 1 we have $R \cdot C - C \cdot R = Q(S,R) = 0$. Thus, we can state that the last condition holds on any Ricci-simple e.c.s. manifold. Finally, we also remark that the last result is an immediate consequence of Theorem 2.2.

7. Conclusions

Let $\overline{M} \times_F \widetilde{N}$ be the warped product of an 1-dimensional manifold $(\overline{M}, \overline{g}), \overline{g}_{11} = \varepsilon = \pm 1$, the warping function $F: \overline{M} \to \mathbb{R}^+$, and an (n-1)-dimensional, $n \ge 4$, semi-Riemannian manifold $(\widetilde{N}, \widetilde{g})$.

If (\tilde{N}, \tilde{g}) is a semi-Riemannian space of constant curvature then $\overline{M} \times_F \tilde{N}$ is a quasi-Eintein conformally flat pseudosymmetric manifold. Evidently, the Friedmann–Lemaître–Robertson–Walker spacetimes belong to this class of manifolds. Furthermore, if the fiber $(\tilde{N}, \tilde{g}), n \ge 5$, is an Einstein manifold, which is not of constant curvature, then $\overline{M} \times_F \tilde{N}$ is a quasi-Einstein nonconformally flat nonpseudosymmetric Ricci-pseudosymmetric manifold. In this case the difference tensor $R \cdot C - C \cdot R$ is expressed by a linear combination of the Tachibana tensors Q(g, R) and Q(S, R) ([8]). If the fibre $(\widetilde{N}, \widetilde{g})$, $n \ge 4$, is a conformally flat Ricci simple manifold such that its scalar curvature $\widetilde{\kappa}$ vanishes then $\overline{M} \times_F \widetilde{N}$ is a non-conformally flat pseudosymmetric manifold, provided that $F = F(x^1) = \exp x^1$ ([10], Proposition 4.2 and Example 4.1). In addition we have $(n-1)(R \cdot C - C \cdot R) = Q(S, C)$ [25].

If the fiber (\tilde{N}, \tilde{g}) , $n \ge 4$, is some Roter-type manifold and the warping function F satisfies (64), then $\overline{M} \times_F \tilde{N}$ is a Roter-type manifold and in consequence a nonconformally flat pseudosymmetric manifold ([36], Theorem 5.1). As was mentioned in Section 2, the tensor $R \cdot C - C \cdot R$ is expressed by a linear combination of some Tachibana tensors.

The above presented facts show that under some conditions imposed on the fiber or the fiber and the warping function of a generalized Robertson–Walker spacetime, such spacetime is a pseudosymmetric or Riccipseudosymmetric manifold and its difference tensor $R \cdot C - C \cdot R$ is expressed by a linear combination of some Tachibana tensors. In this paper we consider an inverse problem. Namely, if the tensors $R \cdot C - C \cdot R$ and Q(S, R) are linearly dependent on a generalized Robertson–Walker spacetime then we determine the warping function, as well as curvature properties of the fiber of such spacetime. In the case where the considered generalized Robertson–Walker spacetimes are 4-dimensional manifolds, it is possible to apply the algebraic classification of spacetimes satisfying some conditions of the pseudosymmetry type given in [27]; see also [39, 42].

Acknowledgments

The authors would like to express their deep gratitude to the referee for his or her valuable comments and suggestions related to the paper.

The first, third, and fifth authors express their thanks to the Institute of Mathematics and Computer Science of the Wrocław University of Technology as well as to the Department of Mathematics of the Wrocław University of Environmental and Life Sciences for hospitality during their stay in Wrocław. The second and fourth authors express their thanks to the Department of Mathematics of Uludağ University in Bursa for hospitality during their stay in Bursa. The authors were supported by grant F-2003/98 of Uludağ University, Bursa, Turkey.

References

- Alías L, Romero A, Sánchez M. Compact spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes. In: Dillen F, editor. Geometry and Topology of Submanifolds VII. River Edge, NJ, USA: World Scientic, 1995, pp. 67–70.
- [2] Alías L, Romero A, Sánchez M. Spacelike hypersurfaces of constant mean curvature and Calabi-Einstein type problems, Tôhoku Math J 1997; 49: 337–345.
- [3] Beem JK, Ehrlich PE. Global Lorentzian Geometry. New York, NY, USA: Marcel Dekker, 1981.
- [4] Beem JK, Ehrlich PE, Powell TG. Warped product manifolds in relativity. In: Rassias TM, editor. Selected Studies: Physics-Astrophysics, Mathematics, History of Sciences, A Volume Dedicated to the Memory of Albert Einstein. Amsterdam, the Netherlands: North-Holland, 1982, pp. 41–56.
- [5] Belkhelfa M, Deszcz R, Głogowska M, Hotloś M, Kowalczyk D, Verstraelen L. On some type of curvature conditions. Banach Center Publ Inst Math Polish Acad Sci 2002; 57: 179–194.
- [6] Besse AL. Einstein Manifolds. Berlin, Germany; Springer-Verlag, 1987.
- [7] Blanco OF, Sánchez M, Senovilla JMM. Structure of second-order symmetric Lorentzian manifolds. J Eur Math Soc 2013; 15: 595–634.

- [8] Chojnacka-Dulas J, Deszcz R, Głogowska M, Prvanović M. On warped product manifolds satisfying some curvature conditions. J Geom Phys 2013; 74: 328–341.
- [9] Defever F, Deszcz R. On semi-Riemannian manifolds satisfying the condition $R \cdot R = Q(S, R)$. In: Verstraelen L, editor. Geometry and Topology of Submanifolds, III. Teaneck, NJ, USA: World Scientific, 1991, pp. 108–130.
- [10] Defever F., Deszcz R., Hotloś M., Kucharski M. and Şentürk Z.: Generalisations of Robertson-Walker spaces. Ann. Univ. Sci. Budapest. Eötvös Sect. Math. 43, 2000, 13-24.
- [11] Defever F, Deszcz R, Prvanović M. On warped product manifolds satisfying some curvature condition of pseudosymmetry type. Bull Greek Math Soc 1994; 36: 43–67.
- [12] Defever F, Deszcz R, Verstraelen L, Vrancken L. On pseudosymmetric space-times. J Math Phys 1994; 35: 5908– 5921.
- [13] Derdziński A, Roter W. Some theorems on conformally symmetric manifolds. Tensor (NS) 1978; 32: 11–23.
- [14] Derdziński A, Roter W. Some properties of conformally symmetric manifolds which are not Ricci-recurrent. Tensor (NS) 1980; 34: 11–20.
- [15] Derdzinski A, Roter W. Projectively flat surfaces, null parallel distributions, and conformally symmetric manifolds. Tohoku Math J 2007; 59: 565–602.
- [16] Derdzinski A, Roter W. On compact manifolds admitting indefinite metrics with parallel Weyl tensor. J Geom Phys 2008; 58: 1137–1147.
- [17] Derdzinski A, Roter W. The local structure of conformally symmetric manifolds. Bull Belg Math Soc Simon Stevin 2009; 16: 117–128.
- [18] Derdzinski A, Roter W. Compact pseudo-Riemannian manifolds with parallel Weyl tensor. Ann Glob Anal Geom 2010; 37: 73–90.
- [19] Deszcz R. On pseudosymmetric spaces. Bull Soc Belg Math Ser A 1992; 44: 1–34.
- [20] Deszcz R. On pseudosymmetric warped product manifolds. In: Dillen F, editor. Geometry and Topology of Submanifolds, V. River Edge, NJ, USA: World Scientific, 1993, pp. 132–146.
- [21] Deszcz R, Głogowska M. Some nonsemisymmetric Ricci-semisymmetric warped product hypersurfaces. Publ Inst Math (Beograd) (NS) 2002; 72: 81–93.
- [22] Deszcz R, Głogowska M, Hotloś M, Sawicz K. A survey on generalized Einstein metric conditions. In: Yau ST, editor. Advances in Lorentzian Geometry: Proceedings of the Lorentzian Geometry Conference in Berlin, AMS/IP Studies in Advanced Mathematics, 2011; 49, 27–46.
- [23] Deszcz R, Głogowska M, Hotloś M, Şentürk Z. On certain quasi-Einstein semisymmetric hypersurfaces. Ann Univ Sci Budapest Eötvös Sect Math 1998; 41: 151–164.
- [24] Deszcz R, Głogowska M, Hotloś M, Verstraelen L. On some generalized Einstein metric conditions on hypersurfaces in semi-Riemannian space forms. Colloq Math 2003; 96: 149–166.
- [25] Deszcz R, Głogowska M, Hotloś M, Zafindratafa G. Hypersurfaces in spaces of constant curvature satisfying some curvature conditions of pseudosymmetric type. In press.
- [26] Deszcz R, Głogowska M, Plaue M, Sawicz K, Scherfner M. On hypersurfaces in space forms satisfying particular curvature conditions of Tachibana type. Kragujevac J Math 2011; 35: 223–247.
- [27] Deszcz R, Haesen S, Verstraelen L. Classification of space-times satisfying some pseudo-symmetry type conditions. Soochow J Math 2004; 30: 339–349.
- [28] Deszcz R, Haesen S, Verstraelen L. On natural symmetries. In: Mihai A, Mihai I, Miron R, editors. Topics in Differential Geometry. Bucharest, Romania: Romanian Academy, 2008, pp. 249–308.
- [29] Deszcz R, Hotloś M. On a certain subclass of pseudosymmetric manifolds. Publ Math Debrecen 1998; 53: 29-48.
- [30] Deszcz R, Hotloś M. On hypersurfaces with type number two in spaces of constant curvature, Ann Univ Sci Budapest Eötvös Sect Math 2003; 46: 19–34.

- [31] Deszcz R, Hotloś M. On some pseudosymmetry type curvature condition. Tsukuba J Math 2003; 27: 13–30.
- [32] Deszcz R, Hotloś M, Şentürk Z. On some family of generalized Einstein metric conditions. Demonstratio Math 2001; 34: 943–954.
- [33] Deszcz R, Hotloś M, Şentürk Z. On curvature properties of certain quasi-Einstein hypersurfaces. Int J Math 2012; 23: 1250073.
- [34] Deszcz R, Kowalczyk D. On some class of pseudosymmetric warped products. Colloq Math 2003; 97: 7–22.
- [35] Deszcz R, Kucharski M. On curvature properties of certain generalized Robertson-Walker spacetimes. Tsukuba J Math 1999; 23: 113–130.
- [36] Deszcz R, Scherfner M. On a particular class of warped products with fibres locally isometric to generalized Cartan hypersurfaces. Colloq Math 2007; 109: 13–29.
- [37] Deszcz R, Verstraelen L, Vrancken L. On the symmetry of warped product spacetimes. Gen Relativ Grav 1991; 23: 671–681.
- [38] Ehrlich PE, Jung YT, Kim SB. Constant scalar curvatures on warped product manifolds. Tsukuba J Math 1996; 20: 239–256.
- [39] Eriksson I, Senovilla JMM. Note on (conformally) semi-symmetric spacetimes. Class Quantum Grav 2010; 27: 027001.
- [40] Głogowska M. On a curvature characterization of Ricci-pseudosymmetric hypersurfaces. Acta Math Scientia 2004; 24 B: 361–375.
- [41] Głogowska M. On quasi-Einstein Cartan type hypersurfaces. J Geom Phys 2008; 58: 599–614.
- [42] Haesen S, Verstraelen L. Classification of the pseudosymmetric space-times. J Math Phys 2004 45: 2343–2346.
- [43] Haesen S, Verstraelen L. Properties of a scalar curvature invariant depending on two planes. Manuscripta Math 2007; 122: 59–72.
- [44] Haesen S, Verstraelen L. Natural intrinsic geometrical symmetries. Symmetry Integrability Geom Meth Appl 2009;
 5: 086.
- [45] Hotloś M. On a certain curvature condition of pseudosymmetry type. To appear.
- [46] Jahanara B, Haesen S, Şentürk Z, Verstraelen L. On the parallel transport of the Ricci curvatures. J Geom Phys 2007; 57: 1771–1777.
- [47] Kowalczyk D. On some subclass of semisymmetric manifolds. Soochow J Math 2001; 27: 445-461.
- [48] Kowalczyk D. On the Reissner-Nordström-de Sitter type spacetimes. Tsukuba J Math 2006; 30: 363–381.
- [49] Kruchkovich GI. On some class of Riemannian spaces. Trudy Sem. po Vekt. i Tenz. Analizu 1961; 11: 103–120 (in Russian).
- [50] Murathan C, Arslan K, Deszcz R, Ezentaş R, Özgür C. On a certain class of hypersurfaces of semi-Euclidean spaces. Publ Math Debrecen 2001; 58: 587–604.
- [51] O'Neill B. Semi-Riemannian Geometry with Applications to Relativity. New York, NY, USA: Academic Press, 1983.
- [52] Özgür C. Hypersurfaces satisfying some curvature conditions in the semi-Euclidean space. Chaos Solitons Fractals 2009; 39: 2457–2464.
- [53] Sánchez M. On the geometry of generalized Robertson-Walker spacetimes: geodesics. Gen Relativ Grav 1998; 30: 915–932.
- [54] Sánchez M. On the geometry of generalized Robertson-Walker spacetimes: curvature and Killing fields. J Geom Phys 1999; 31: 1–15.
- [55] Senovilla JMM. Second-order symmetric Lorentzian manifolds: I. Characterization and general results. Class Quantum Grav 2008; 25: 245011.
- [56] Shaikh AA, Kim YH, Hui SK. On Lorentzian quasi-Einstein manifolds. J Korean Math Soc 2011; 48: 669–689.

ARSLAN et al./Turk J Math

- [57] Stephani H, Kramer D, MacCallum M, Hoenselaers C, Herlt E. Exact Solutions of the Einstein's Field Equations. Cambridge Monographs on Mathematical Physics. 2nd ed. Cambridge, UK: Cambridge University Press, 2003.
- [58] Szabó ZI. Structure theorems on Riemannian spaces satisfying $R(X, Y) \cdot R = 0$. I. The local version. J Differential Geom 1982; 17: 531–582.
- [59] Szabó ZI. Structure theorems on Riemannian spaces satisfying $R(X, Y) \cdot R = 0$. II. The global version. Geom Dedicata 1985; 19: 65–108.