

1-1-2010

## Effects of moist chilling, gibberellic acid, and scarification on seed dormancy in the rare endemic *Pedicularis olympica* (Scrophulariaceae)

SERAP KIRMIZI

GÜRCAN GÜLERYÜZ

HÜLYA ARSLAN

FATMA SELCEN SAKAR

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

---

### Recommended Citation

KIRMIZI, SERAP; GÜLERYÜZ, GÜRCAN; ARSLAN, HÜLYA; and SAKAR, FATMA SELCEN (2010) "Effects of moist chilling, gibberellic acid, and scarification on seed dormancy in the rare endemic *Pedicularis olympica* (Scrophulariaceae)," *Turkish Journal of Botany*. Vol. 34: No. 3, Article 8. <https://doi.org/10.3906/bot-0912-287>

Available at: <https://journals.tubitak.gov.tr/botany/vol34/iss3/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Effects of moist chilling, gibberellic acid, and scarification on seed dormancy in the rare endemic *Pedicularis olympica* (Scrophulariaceae)

Serap KIRMIZI<sup>1</sup>, Gürcan GÜLERYÜZ<sup>2,\*</sup>, Hülya ARSLAN<sup>2</sup>, F. Selcen SAKAR<sup>2</sup>

<sup>1</sup>Gemlik Asım Kocabıyık Graduate Vocational School, Horticulture Programme, Uludağ University, 16600, Gemlik, Bursa - TURKEY

<sup>2</sup>Faculty of Science and Arts, Department of Biology, Uludağ University, 16059, Görükle, Bursa - TURKEY

Received: 25.12.2009

Accepted: 16.02.2010

**Abstract:** We investigated the germination requirements of a rare endemic plant species, *Pedicularis olympica* Boiss. (Scrophulariaceae), which grows in wet habitats on Uludağ Mountain, Turkey. We studied the effects of scarification, moist chilling (+4 °C) for 15 days, different doses of gibberellic acid (GA<sub>3</sub>; 100, 150, and 250 ppm) and combined hormone and moist chilling treatments under dark (20 °C) and photoperiod (20/10 °C; 12/12 h, respectively) conditions. GA<sub>3</sub> was able to break dormancy. The highest germination rate was found in the seeds treated with 250 ppm GA<sub>3</sub>; 64% of these seeds germinated when treated with moist chilling and incubated in the dark, while 75% germinated under photoperiod conditions. Moreover, mean germination times were significantly lower in the seeds treated with 250 ppm GA<sub>3</sub>. Significant differences in final germination percentages and mean germination times were found only for treatments under photoperiod conditions. Scarification also permitted germination; the highest germination rate (78%) was found with 15 min of scarification. Dormancy was not broken by moist chilling for up to 90 days. The germination requirements of dormant *P. olympica* seeds are found to be consistent with characteristics of its habitat. This can be considered an ecological advantage for the species' establishment and persistence.

**Key words:** Seed dormancy, *Pedicularis olympica*, endemic, stratification, gibberellic acid, scarification

### Nadir ve endemik *Pedicularis olympica* (Scrophulariaceae) türünde nemli soğuklama, gibberellik asit ve skarifikasyonun tohum dormansisi üzerindeki etkileri

**Özet:** Bu çalışmada Uludağ'da nemli alanlarda yayılış gösteren nadir endemik *Pedicularis olympica* Boiss. (Scrophulariaceae) türünün çimlenme gereksinimleri araştırılmıştır. Çalışmada, skarifikasyon, 15 gün nemli soğuklama (+4 °C) ve farklı dozlarda gibberellik asit (GA<sub>3</sub>; 100, 150 ve 250 ppm) ile hormon ve nemli üşütme kombinasyonlarının karanlık (20 °C) ve fotoperiyot (sırasıyla 20/10 °C; 12/12 s) şartlarında etkileri araştırılmıştır. GA<sub>3</sub> uygulamalarının dormansiyi kırmada etkili olduğu saptanmıştır. En yüksek çimlenme oranı 250 ppm GA<sub>3</sub> ile muamele edilmiş tohumlarda bulunmuştur; karanlık şartlarda nemli üşütmede % 64 iken fotoperiyotta % 75 çimlenme gerçekleşmiştir. Ayrıca, 250 ppm GA<sub>3</sub> ile muamele edilen tohumlarda ortalama çimlenme süresi de anlamlı derecede kısalmıştır. Final çimlenme oranları

\* E-mail: gurcan@uludag.edu.tr

ve ortalama çimlenme süreleri arasında anlamlı fark grupları sadece fotoperiyot şartları altında bulunmuştur. Skarifikasyon çimlenmeyi uyarmıştır ve en yüksek çimlenme 15 dakika skarifikasyon sonucunda bulunmuştur (% 78). 90 güne kadar yapılan nemli üşütme muamelelerinde dormansi kırılmamıştır. *P. olympica* türünün çimlenme gereksinimlerinin bulunduğu habitatla uyum içinde olduğu bulunmuştur. Bu da türün yerleşme ve devamlılığında bir ekolojik avantaj olarak değerlendirilebilir.

**Anahtar sözcükler:** Tohum dormansisi, *Pedicularis olympica*, endemik, soğuklama, gibberellik asit, skarifikasyon

## Introduction

Seedling establishment is a critical stage in the life history of any plant species that relies on sexual reproduction for the persistence of its populations (Grubb, 1977; Harper, 1979; Bu et al., 2008). Variations in seed dispersal efficacy or germination percentage are often interpreted as reflecting adaptations to specific ecological conditions (Grime et al., 1981; Nishitani & Masuzawa, 1996). A large number of endemic species are found in alpine habitats (Väre et al., 2003), and seeds of such species must possess elaborate dormancy mechanisms to survive the harsh climatic conditions that ensue immediately after their seed maturation and dispersal (Billings & Mooney, 1968). Previous studies on alpine endemic species suggest that no universal alpine germination strategy exists (Gimenez-Benavides et al., 2005), but these species generally exhibit innate or enforced dormancy that prevents germination during the unfavourable conditions of winter (Körner, 1999).

Dormancy can be defined in multiple ways. Seed dormancy can be defined simply as inhibited germination of an intact viable seed to optimise the distribution of germination over time (Bewley & Black, 1983; Hilhorst, 1995). This inhibition of germination has evolved differently across species for adaptation to the prevailing environment so that germination occurs when conditions are likely to be suitable for establishment of a new generation (Finch-Savage & Leubner-Metzger, 2006). Baskin and Baskin (1998) have reported that among 300 temperate herbaceous species, dormancy breaking and germination requirements are not phylogenetically constrained. According to the classification system proposed by Baskin and Baskin (1998, 2004), physiological dormancy is the most common form, found in seeds of gymnosperms and of all angiosperm clades. Physiological dormancy is the most prevalent

form of dormancy in temperate seed banks. It may be deep, intermediate, or non-deep (Finch-Savage & Leubner-Metzger, 2006). Dormancy may be strong to weak, and the extent of dormancy present at any particular moment is referred to as the degree of dormancy. Dormancy patterns are similar for closely related taxa but may differ within a family, even between co-occurring species with similar life histories (Karlsson et al., 2006). The germination requirements may represent extreme conditions that do not normally exist, thus inhibiting complete germination.

Soil moisture and soil temperature are the most important factors regulating seed behaviour under natural conditions (Bewley & Black, 1982). In some wetland species, germination is not limited by secondary dormancy; once seeds have come out of primary dormancy, germination can occur at any time during the growing season if conditions are favourable (Baskin & Baskin, 1993a, 1993b; Milberg, 1994; Jensen, 2004). It has been shown that seed germination in wet meadows is negatively affected by the thickness of the litter layer (Facelli & Pickett, 1991; Diemer et al., 2001).

Wetland species require high temperatures for germination and respond to large daily temperature fluctuations (Grime et al., 1981). These characteristics are regarded as adaptations to delay germination until the water table recedes and the soil surface becomes exposed in the spring, thus creating oxygenated conditions with warmer soil temperatures and larger temperature fluctuations (Grime et al., 1981; Thompson & Grime, 1983). Many studies have examined the influence of various treatments, including gibberellins (Li et al, 2007; Ren & Guan, 2008) and stratification (Schütz & Rave, 1999; Brändel & Jensen, 2005; Brändel, 2006; Kettenring & Galatowisch, 2007) on seed germination in wetlands.

These studies have pointed out the important role of dormancy in the reproductive success of wetland plants.

Germination requirements for native species, particularly for rare and/or endemic species, are important in conservation biology (Cerabolini et al., 2004). Investigation of germination requirements may show how a species' germination process is adapted to habitat conditions, how it is regulated by environmental factors (Van Assche et al., 2002), and how it influences subsequent seedling establishment in a particular habitat (Schütz & Rave, 1999). *Pedicularis olympica* Boiss. (Scrophulariaceae) is an endemic species from Uludağ Mountain, Turkey. The species' habitat has been damaged by heavy recreational and winter sports activity. The status of the species is Vulnerable (VU) according to the IUCN classification (Güleryüz, 1998). The germination requirements of the species were not previously studied. In this study, we evaluated the germination requirements of *P. olympica* by testing its responses to GA<sub>3</sub>, moist chilling, and scarification treatments in relation to the breaking of dormancy.

## Materials and methods

### Study area

Uludağ is the highest mountain in the Marmara region, including Thrace and the north-western side of the Anatolian peninsula. The climate of the mountain varies with elevation, resulting in high biological diversity. Because of its high plant diversity, Uludağ Mountain is one of the important plant areas (IPAs) in Turkey (Güleryüz et al., 2005). To preserve the unique plant communities and various geomorphological features, an area of 11,338 ha was designated as a National Park in 1961. This area was enlarged to 12,762 ha in 1998 (Arslan et al., 1999). A Mediterranean-type climate is found at lower elevations, near the city of Bursa on the north-western side of the mountain. Rainy, partially mild, micro-thermic climates with icy winters are found at higher altitudes (Güleryüz, 1992).

### Study species and habitat characteristics

*Pedicularis olympica* (Scrophulariaceae) is a perennial herb with erect, unbranched, eglandular-

pubescent, or almost glabrous stems. Its leaves are linear-oblong, simply pinnatisect or pinnatifid, and crenate-serrate. The basal leaves are petiolate; the cauline leaves are several and alternate. The inflorescence is a many-flowered, oblong, sublanate spike that elongates in fruit. In general, seeds of *Pedicularis* species are very small, containing a minimal embryo with a linear shape, and are protected by a thin brown-coloured seed coat. The seeds are laterally compressed and elliptic in outline. The seed surface is reticulate, and the embryo is linear and axile (Ellis, 1985).

*P. olympica* is known only from Uludağ Mountain, where it is found in wet habitats and on stream banks, especially near springs. The species grows on granite substrates. It is one of the first species to flower in spring after snow melts in the subalpine zone, forming an attractive landscape of rose-red flowers in meadows. The flowering period is from May to June (Davis, 1975; Güleryüz, 1998).

### Germination tests

Mature seeds of *P. olympica* were collected from the subalpine belt of Uludağ Mountain between 1800 and 1850 m in elevation during June 2008. Germination experiments were started immediately after drying and cleaning the seeds. Sterile plastic 9-mm petri dishes were used for germination experiments. Seeds were surface-sterilised for 3 min with 5% sodium hypochlorite and then rinsed with tap water. Some preliminary treatments were made to determine the moist chilling duration and GA<sub>3</sub> doses. These were 30, 60, and 90 days of moist chilling with 250, 500, and 1000 ppm GA<sub>3</sub>, and control without GA<sub>3</sub> (distilled water). After the preliminary treatments the test solutions were chosen as 100, 150, and 250 ppm GA<sub>3</sub> and distilled water as a control. All of the hormone solutions were analytical grade. The hormone solutions were applied as a pre-treatment for 24 h of imbibition, after which the seeds were rinsed with distilled water. Four replicates of 25 seeds per petri dish were made. Half of the replicates were incubated under photoperiod conditions: 20/10 °C (12/12 h). The other half of the replicates were incubated at 20 °C in the dark. For dark treatments, the petri dishes were wrapped with aluminium foil. Moist chilling was achieved by incubating seeds under wet and cold (+4 °C) conditions for 15 days. The

number of germinated seeds was counted and the germinated seeds were removed every day for up to 25 days. Seeds were recorded as having germinated when the radicle emerged from the testa. Mean germination time (MGT) was calculated from the germination counts and used to determine the speed of germination. Final germination percentages and mean germination times were determined at the end of the 25-day incubation period. For scarification, seeds were treated with 80% sulphuric acid and then rinsed with tap water.

### Statistical Analyses

Data were analysed using one-way analysis of variance (ANOVA). Differences among treatment means were tested by Tukey's honest significance difference (HSD) test. All tests were performed at the significance level of  $\alpha = 0.05$  using the Statistica 6.0 (StatSoft, 1984-1995) software package.

### Results and discussion

Results for pre-treatments with GA<sub>3</sub> and moist chilling for seeds of *P. olympica* under photoperiod conditions are given in Table 1. Seeds failed to germinate in control group, in which neither GA<sub>3</sub> nor moist chilling was applied. Seeds also failed to germinate in control moist chilling treatments with distilled water, except after 60 and 90 days of chilling,

which produced 8.3% and 7.8% germination, respectively (Table 1). The highest germination rate after 21 days of moist chilling was found in seeds treated with 250 ppm GA<sub>3</sub> (93.2%). However, germination rate decreased with increasing GA<sub>3</sub> above 250 ppm. Ren and Guan (2008) have found similar results for different *Pedicularis* species; longer durations of stratification with GA<sub>3</sub> lowered germination. Mean germination times decreased in moist-chilled compared to non-chilled ones. After the preliminary treatments, GA<sub>3</sub> doses of 100, 150, and 250 ppm and combinations of GA<sub>3</sub> with 15 days of moist chilling were chosen as treatments for the main experiment.

Seeds failed to germinate in control treatments with distilled water under both dark and photoperiod conditions (Figure). There was a clear difference among GA<sub>3</sub> doses of 100, 150, and 250 ppm. Germination increased with increasing GA<sub>3</sub> dose under both dark and photoperiod conditions. The highest germination rates were found in the seeds treated with 250 ppm GA<sub>3</sub>: 64% for dark conditions with moist chilling and 75% for photoperiod conditions (Figure). Final germination percentages and MGTs are presented in Table 2. Significant differences in the final germination percentages and MGTs were found only for treatments under photoperiod conditions ( $P < 0.005$ ).

Table 1. Preliminary treatments results for determination of GA<sub>3</sub> doses and duration of moist chilling on seeds of *Pedicularis olympica* (Mean  $\pm$  Standard Error, nd: not determined).

Moist Chilling Duration (days)	GA <sub>3</sub> concentration (ppm)			
	0	250	500	1000
Final Germination (%)				
0 (control)	0 $\pm$ 0	52.0 $\pm$ 6.7	96.8 $\pm$ 1.8	97.6 $\pm$ 3.3
21	0 $\pm$ 0	93.2 $\pm$ 1.8	88.4 $\pm$ 3.8	87.2 $\pm$ 7.8
30	0 $\pm$ 0	nd	nd	nd
60	8.8 $\pm$ 5.6	nd	nd	nd
90	7.3 $\pm$ 1.9	nd	nd	nd
Mean germination time (days)				
0 (control)	0 $\pm$ 0	16.0 $\pm$ 0.8	7.4 $\pm$ 0.8	7.2 $\pm$ 0.1
21	0 $\pm$ 0	4.4 $\pm$ 0.3	2.5 $\pm$ 0.8	2.3 $\pm$ 0.4
30	0 $\pm$ 0	nd	nd	nd
60	9.4 $\pm$ 0.5	nd	nd	nd
90	7.4 $\pm$ 2.1	nd	nd	nd






Figure. Cumulative germination percentage diagrams for *Pedicularis olympica* seeds in 4 different treatments under dark (20 °C) and photoperiod (20/10 °C; 12/12 h, respectively) conditions.

Scarification with sulphuric acid promoted germination (Table 2). Since the scarification with sand paper would be destructive for small seeds, we treated the seeds with 80% sulphuric acid for 5, 10, or 15 min. The highest germination rate was found with 15 min of scarification (78%). Scarification was also found to be the most effective method for promoting germination in 7 of 8 *Pedicularis* species from Yunnan, China (Li et al., 2007). The authors of that study reported that the germination behaviours of the 8 *Pedicularis* species varied considerably. It is thought that wetland species usually require light for germination (Grime et al., 1981). However, our results suggest that *P. olympica* does not require light for germination. This may be due to its habitat; *P. olympica* prefers to grow near stream banks, especially near springs.

According to the results found in this study, *P. olympica* species probably exhibits a combination of

physical and physiological dormancy (PY+PD).  $GA_3$  was able to break dormancy; scarification also permitted germination.

The germination rate resulting from 15 min of scarification was similar to the results of 250 ppm  $GA_3$  with moist chilling and photoperiod germination conditions. While MGT was 16 days for seeds treated with 250 ppm  $GA_3$  and moist chilling that were germinated under photoperiod conditions, it decreased to 8 days with scarification. *Pedicularis* species do not have thick seed coats, but endosperm can act as a mechanical inhibitor of seed germination in several angiosperms (Finch-Savage & Leubner-Metzger, 2006). A decline in the mechanical resistance of the micropylar endosperm may be a prerequisite for radicle protrusion during seed germination (Hilhorst, 1995; Leubner-Metzger, 2003; Kucera et al., 2005). The weakening of endosperm can be promoted by gibberellic acid and is inhibited (at least in part) by

Table 2. Final germination (%) and mean germination times (MGT; days) of *P. olympica* seeds for different treatment series. [Values are means followed by standard error (n = 4); the values followed by the same letter were not statistically significant at the  $\alpha = 0.05$  level].

Treatment Series		Germination (%)	MGT	
Dark (20 °C)	Control	0	0	
	100 ppm GA <sub>3</sub>	49 <sup>a</sup> ± 3.5	14.2 <sup>a</sup> ± 1.0	
	150 ppm GA <sub>3</sub>	59 <sup>a</sup> ± 3.0	14.0 <sup>a</sup> ± 0.2	
	250 ppm GA <sub>3</sub>	60 <sup>a</sup> ± 4.9	12.7 <sup>a</sup> ± 0.9	
	Stratification 15 days (+4 °C)	Control	0	0
		100 ppm GA <sub>3</sub>	48 <sup>a</sup> ± 8.6	13.7 <sup>a</sup> ± 0.5
		150 ppm GA <sub>3</sub>	60 <sup>a</sup> ± 4.8	14.2 <sup>a</sup> ± 0.8
		250 ppm GA <sub>3</sub>	64 <sup>a</sup> ± 3.3	12.9 <sup>a</sup> ± 0.9
Fotoperiod (20/10 °C; 12/12 h)	Control	0	0	
	100 ppm GA <sub>3</sub>	49 <sup>a</sup> ± 2.5	16.0 <sup>a</sup> ± 0.3	
	150 ppm GA <sub>3</sub>	48 <sup>a</sup> ± 2.8	15.4 <sup>ab</sup> ± 0.5	
	250 ppm GA <sub>3</sub>	52 <sup>a</sup> ± 4.7	16.2 <sup>a</sup> ± 0.4	
	Stratification 15 days (+4 °C)	Control	0	0
		100 ppm GA <sub>3</sub>	44 <sup>b</sup> ± 8.2	9.4 <sup>bc</sup> ± 1.8
		150 ppm GA <sub>3</sub>	48 <sup>b</sup> ± 9.2	11.9 <sup>b</sup> ± 7.7
		250 ppm GA <sub>3</sub>	75 <sup>a</sup> ± 4.1	10.6 <sup>bc</sup> ± 0.7
Scarification (80% H <sub>2</sub> SO <sub>4</sub> )	5 min	26 <sup>c</sup> ± 2.6	2.3 <sup>b</sup> ± 0.1	
	10 min	47 <sup>b</sup> ± 5.5	2.5 <sup>b</sup> ± 0.1	
	15 min	78 <sup>a</sup> ± 2.5	3.4 <sup>a±</sup> 0.2	

abscisic acid (ABA) (Finch-Savage & Leubner-Metzger, 2006). The germination of tomato is inhibited by ABA, but surgical removal of the micropylar cap permits germination even in the presence of ABA (Liptay & Schopfer, 1983). Endosperm weakening is also known to be required for radicle protrusion in Solanaceae species (Finch-Savage & Leubner-Metzger, 2006). GA<sub>3</sub> might enhance the growth potential of the embryo (Karssen & Lacka, 1986; Karssen et al., 1989) or induce degradation of food reserves in endosperm endosperm by stimulating hydrolytic enzyme activity (Da Silva et al., 2005) of *P. olympica* seeds. In nature, some of the ungerminated seeds may remain alive and serve as a soil seed bank, waiting for more favourable conditions. This strategy may be common among high-elevation wetland plants (Leck & Simpson, 1987; Kaye, 1997). Li et al. (2007) reported that GA<sub>3</sub>, scarification and surface sowing promoted germination in the 8 *Pedicularis* species they studied and suggested that allowing sufficient time for

germination may be useful for germination under natural conditions.

Alpine and subalpine plants provide good opportunities to determine differences between species in germination responses. Investigations of the germination ecology of alpine plants are rare; therefore, the factors and mechanisms that regulate germination in alpine habitats are poorly known (Baskin & Baskin, 1998). *P. olympica* seeds exhibit dormancy characteristics that are consistent with environmental conditions in its habitat. This can be considered an ecological advantage for the species' establishment and persistence.

The attractive rare endemic plant *Pedicularis olympica* is vulnerable to habitat degradation on Uludağ Mountain, its only known locality. It spreads in wet meadows within the subalpine belt of Uludağ Mountain and has very attractive rose red flowers that can be used for ornamental purposes. Conservation biologists must develop new strategies for protecting

species and ecosystems because of habitat losses, landscape alteration, and extinction of species, communities and even ecosystems (Hamilton, 1994). Ex situ conservation from seed is regarded as a viable and inexpensive method for threatened plant species, but the germination requirements of alpine species are poorly known, with a few exceptions (Cerabolini et al., 2004; Gimenez-Benavides et al., 2005). The

germination requirements found in our study will be useful for future ex situ conservation of *P. olympica*.

## References

- Arslan H, Güleriyüz G, Gökçeoğlu M & Rehder H (1999). Vegetation mosaic around the second center of tourism development in the Uludağ Mountain, Bursa, Turkey. *Turk J Bot* 23: 233-239.
- Baskin CC & Baskin JM (1998). *Seeds. Ecology, Biogeography and Evolution of Dormancy and Germination*. San Diego: Academic Press.
- Baskin CC, Baskin JM & Chester, EW (1993a). Seed germination ecophysiology of four summer annual mudflat species of *Cyperaceae*. *Aquat Bot* 45: 41-52.
- Baskin CC, Baskin JM & Leck MA (1993b). After ripening pattern during cold stratification of achenes of perennial *Asteraceae* from Eastern North America and evolutionary implications. *Plant Spec Biol* 8: 61-65.
- Baskin JM & Baskin CC (2004). A classification system for seed dormancy. *Seed Sci Res* 14: 1-16.
- Bewley JD & Black M (1983). *Physiology and Biochemistry of Seeds In Relation to Germination. Vol 1. Development, germination, and growth*, pp. 81-100. Berlin, Heidelberg, New York: Springer Verlag.
- Bewley JD & Black M (1982). *Physiology and Biochemistry of Seeds In Relation to Germination, Vol. 2. Viability, Dormancy and Environmental Control*, pp. 212-223. Berlin, Heidelberg, New York: Springer Verlag.
- Billings WD & Mooney HA (1968). The ecology of arctic and alpine plants. *Biol Rev* 43: 481-529.
- Brändel M & Jernsen K (2005). Effect of temperature on dormancy and germination of *Eupatorium cannabinum* L. achenes. *Seed Sci Res* 15: 143-151.
- Brändel M (2006). Effect of temperatures on dormancy and germination in three species in the Lamiaceae occurring in northern wetlands. *Wetlands Ecol Man* 14: 11-28.
- Bu H, Du G, Chen X, Xu X, Liu K & Wen S (2008). Community wide germination strategies in alpine meadow on the eastern Qinghai-Tibet plateau: phylogenetic and life history correlates. *Plant Ecol* 195: 87-98.
- Cerabolini B, De Andreis R, Ceriani RM, Pierce S & Raimondi B (2004). Seed germination and conservation of endangered species from the Italian Alps: *Physoplexis comosa* and *Primula glaucescens*. *Biol Conserv* 117: 351-356.
- Da Silva EAA, Toorop PE, Jaap Nijse J, Bewley JD & Hilhorst HWM (2005). Exogenous gibberellins inhibit coffee (*Coffea arabica* cv. Rubi) seed germination and cause cell death in the embryo. *J Exp Bot* 56: 1029-1038.
- Davis PH (Ed.) (1975). *Flora of Turkey and East Aegean Islands*, Vol. 5: p. 772. Edinburgh: Edinburgh University Press.
- Diemer M, Oetiker K & Billeter R (2001). Abandonment alters community composition and canopy structure of Swiss calcareous fens. *Appl Veg Sci* 4: 237-246.
- Ellis RH, Hong TD & Roberts EH (1985). *Handbook for Seed Technology of Genebanks. Vol 2. Compendium of Specific Germination Information and Test Recommendations*, Rome: International Board for Plant Genetic Resources.
- Facelli JM & Pickett STA (1991). Plant litter: its dynamics and effects on plant community structure. *Bot Rev* 57: 1-32.
- Finch-Savage WE & Leubner-Metzger G (2006). Seed dormancy and the control of germination. *New Phytol* 171: 501-523.
- Gimenez-Benavides L, Escudero A & Perez-Garcia F (2005). Seed germination of high mountain Mediterranean species: altitudinal, interpopulation and interannual variability. *Ecol Res* 20: 433-444.
- Grime JP, Mason G, Curtis AV, Rodman J & Band SR (1981). A comparative study of germination characteristics in a local flora. *J Ecol* 69: 1017-1059.
- Grubb PJ (1977). The maintenance of species-richness in plant communities: the importance of the regeneration niche. *Biol Rev* 52: 107-145.
- Güleriyüz G (1998). *Pedicularis olympica* Boiss. (Bit otu) *Scrophulariaceae* (Yüksükotugiller). *The Karaca Arboretum Magazine* 4: 189-190.
- Güleriyüz G (1992). The studies on the nutrient turnover and productivity at some plant communities of the Uludağ Alpine Region. Doctoral dissertation, Graduate School of Natural and Applied Sciences, University of Uludağ, Bursa, Turkey.
- Güleriyüz G, Malyer H, Kaynak G & Özhatay N (2005). Uludağ A2 (A) Bursa. In: Özhatay N, Bayfield A & Atay S (eds.) *Important Plant Areas in Turkey*, İstanbul: WWF Turkey.



- Hamilton MB (1994). Ex situ conservation of wild plant species: time to reassess the genetic assumptions and implications of seed banks. *Conserv Biol* 8: 39-49.
- Harper JL (1979). *Population Biology of Plants*. London, New York: Academic Press.
- Hilhorst HWM (1995). A critical update on seed dormancy. I. Primary dormancy. *Seed Sci Res* 5: 61-73.
- Jensen K (2004). Dormancy patterns, germination ecology and seed-bank types of 20 temperate fen grassland species. *Wetlands* 24: 152-166.
- Karlsson LM, Ericsson J & Milberg P (2006). Seed dormancy and germination in the summer annuals *Galeopsis speciosa*. *Weed Res* 46: 353-361.
- Karssen CM & Lacka E (1986). A revision of the hormone balance theory of seed dormancy: studies on gibberellin and/or abscisic acid-deficient mutants of *Arabidopsis thaliana*. In: Bopp M (ed.) *Plant Growth Substances*, pp. 315-323. Berlin: Springer-Verlag.
- Karssen CM, Zagorski S, Kepczynski J & Groot SPC (1989). A key role for endogenous gibberellins in the control of seed germination. *Ann Bot* 63: 71-80.
- Kaye TN (1997). Seed dormancy in high elevation plants: Implications for ecology and restoration. In: Kaye TN, Liston A, Love RM, Luoma DL, Meinke RJ & Wilson MV (eds.) *Conservation and Management of Native Plants and Fungi*, pp. 115-120. Corvallis, Oregon: Native Plant Society of Oregon.
- Kettenring K & Galatowisch SM (2007). Temperature requirements for dormancy break and seed germination vary greatly among 14 wetland *Carex* species. *Aquat Bot* 87: 209-220.
- Körner C (1999). *Alpine Plant Life*. 2nd edition. Berlin: Springer Verlag.
- Kucera B, Cohn MA & Leubner-Metzger G (2005). Plant hormone interactions during seed dormancy release and germination. *Seed Sci Res* 15: 281-307.
- Leck MA & Simpson RL (1987). Seed Bank of a Freshwater Tidal Wetland: Turnover and Relationship to Vegetation Change. *Am J Bot* 74: 360-370.
- Leubner-Metzger G (2003). Functions and regulation of  $\beta$ -1,3 glucanase during seed germination, dormancy release and after ripening. *Seed Sci Res* 13: 17-34.
- Li AR, Guan KY & Probert RJ (2007). Effects of light, scarification, and gibberellic acid on seed Germination of eight *Pedicularis* species from Yunnan, China. *Hortsci* 42: 1259-1262.
- Liptay A & Schopfer P (1983). Effect of water stress, seed coat restraint, and abscisic acid upon different germination capabilities of two tomato lines at low temperature. *Plant Physiol* 73: 935-938.
- Milberg P (1994). Annual dark dormancy cycle in buried seeds of *Lychnis flos-cuculi*. *Ann Bot Fenn* 31: 163-167.
- Nishitani S & Masuzawa T (1996). Germination characteristics of two species of *Polygonum* in relation to their altitudinal distribution on Mt Fuji, Japan. *Arc Alp Res* 28: 104-110.
- Ren YQ & Guan KY (2008). Effects of moist chilling and GA<sub>3</sub> applications on seed germination of three *Pedicularis* species from Yunan, China. *Seed Sci Technol* 36: 225-229.
- SAS Institute, SAS/STATISTICA Version 6.0 SAS Institute Inc., Cary, North Carolina, USA. 1984-1995.
- Schütz W & Rave G (1999). The effect of cold stratification and light on the seed germination of temperature sedges (*Carex*) from various habitats and implications for regenerative strategies. *Plant Ecol* 144: 215-230.
- Thompson K & Grime JP (1983). A comparative study of germination responses to diurnally-fluctuating temperatures. *J Appl Ecol* 20: 141-156.
- Van Assche J, Van Nerum D & Darius P (2002). The comparative germination ecology of nine *Rumex* species. *Plant Ecol* 159: 131-142.
- Väre H., Lampinen C, Humpries C & Williams P (2003). Taxonomic Diversity of Vascular Plants in the European Alpine Areas. In: Nagy L, Grabherr G, Körner C & Thompson DBA (eds.) *Alpine Biodiversity in Europe*, pp. 133-148. Ecological studies, V 167. Berlin Heidelberg New York: Springer.