human reproduction

CORRIGENDUM

Which is worse? A comparison of ART outcome between women with primary or recurrent endometriomas

Baris Ata¹, Sezcan Mumusoglu², Kiper Aslan³, Ayse Seyhan⁴, Isil Kasapoglu³, Berrin Avci³, Bulent Urman¹, Gurkan Bozdag², and Gurkan Uncu^{3,*}

Department of Obstetrics and Gynecology, Koc University School of Medicine, Davutpasa Caddesi No 4, Topkapi, Istanbul 34010, Turkey ²Department of Obstetrics and Gynecaology, Hacettepe University School of Medicine, Sihhiye, Ankara 06100, Turkey ³Department of Obstetrics and Gynecology, Uludag University School of Medicine, Gorukle, Bursa 16059, Turkey ⁴Women's Health and Assisted Reproduction Center of the American Hospital of Istanbul, Guzelbahce Sokak, Nisantasi, Istanbul 34365, Turkey

*Correspondence address. Division of Reproductive Endocrinology and Infertility, Department of Obstetrics and Gynecology, Uludag University School of Medicine, Gorukle, Bursa, Turkey. Tel: +90-224-295-0000; E-mail: guncu@gurkanuncu.com

Hum Reprod 2017;32: doi:10.1093/humrep/dex099

The authors would like to apologize for errors in the affiliations listed for four of the authors (Isil Kasapoglu, Berrin Avci, Bulent Urman and Gurkan Uncu) of the above article. The correct affiliations are:

Isil Kasapoglu

³Department of Obstetrics and Gynecology, Uludag University School of Medicine, Gorukle, Bursa 16059, Turkey

Berrin Avci

³Department of Obstetrics and Gynecology, Uludag University School of Medicine, Gorukle, Bursa 16059, Turkey

Bulent Urman

Department of Obstetrics and Gynecology, Koc University School of Medicine, Davutpasa Caddesi No 4, Topkapi, Istanbul 34010, Turkey

Gurkan Uncu

³Department of Obstetrics and Gynecology, Uludag University School of Medicine, Gorukle, Bursa 16059, Turkey

The electronic version of this article has been updated at https://doi. org/10.1093/humrep/dex099. The authors would like to assure readers that this does not affect any content of the article.