

1-1-2018

On the Chebyshev coefficients for a general subclass of univalentfunctions

ŞAHSENE ALTINKAYA

SİBEL YALÇIN TOKGÖZ

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ALTINKAYA, ŞAHSENE and TOKGÖZ, SİBEL YALÇIN (2018) "On the Chebyshev coefficients for a general subclass of univalentfunctions," *Turkish Journal of Mathematics*: Vol. 42: No. 6, Article 3. <https://doi.org/10.3906/mat-1510-53>

Available at: <https://journals.tubitak.gov.tr/math/vol42/iss6/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On the Chebyshev coefficients for a general subclass of univalent functions

Şahsene ALTINKAYA*, Sibel YALÇIN

Department of Mathematics, Faculty of Arts and Science, Bursa Uludağ University, Bursa, Turkey

Received: 15.10.2015

Accepted/Published Online: 03.03.2017

Final Version: 27.11.2018

Abstract: In this work, considering a general subclass of univalent functions and using the Chebyshev polynomials, we obtain coefficient expansions for functions in this class.

Key words: Chebyshev polynomials, analytic and univalent functions, coefficient bounds, subordination

1. Introduction and definitions

Let D be the open unit disk $\{z \in \mathbb{C} : |z| < 1\}$ and A be the class of functions analytic in D , satisfying the conditions

$$f(0) = 0 \text{ and } f'(0) = 1.$$

Then each function f in A has the following Taylor expansion

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n. \quad (1)$$

Furthermore, by S we shall denote the class of all functions in A that are univalent in D .

If the functions f and g are analytic in D , then f is said to be subordinate to g , written as

$$f(z) \prec g(z), \quad (z \in D),$$

if there exists a Schwarz function $w(z)$, analytic in D , with

$$w(0) = 0 \text{ and } |w(z)| < 1 \quad (z \in D)$$

such that

$$f(z) = g(w(z)) \quad (z \in D).$$

Denote by K the subclass of S of convex functions so that $f \in K$ if and only if, for $z \in D$,

$$\Re \left(1 + \frac{zf''(z)}{f'(z)} \right) > 0.$$

The Fekete–Szegő functional $|a_3 - \mu a_2^2|$ for normalized univalent functions of the form given by (1) is well known for its rich history in geometric function theory. Its origin was in the disproof by Fekete and Szegő

*Correspondence: sahsene@uludag.edu.tr

2010 AMS Mathematics Subject Classification: 30C45, 30C50

of the 1933 conjecture of Littlewood and Paley that the coefficients of odd univalent functions are bounded by unity (see [5]). The functional has since received great attention, particularly in many subclasses of the family of univalent functions. Nowadays, it seems that this topic had become of interest among researchers (see, for example, [1, 2, 7, 8]).

Chebyshev polynomials have become increasingly important in numerical analysis, from both theoretical and practical points of view. There are four kinds of Chebyshev polynomials. The majority of books and research papers dealing with specific orthogonal polynomials of the Chebyshev family contain mainly results of Chebyshev polynomials of the first and second kinds $T_n(x)$, $U_n(x)$ and their numerous uses in different applications; see, for example, Doha [3] and Mason [6].

The Chebyshev polynomials of the first and second kinds are well known. In the case of a real variable x on $(-1, 1)$, they are defined by

$$T_n(x) = \cos n\theta,$$

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta},$$

where the subscript n denotes the polynomial degree and where $x = \cos \theta$.

Definition 1 A function $f \in A$ is said to be in the class $L(\alpha, t)$, $\alpha \geq 0$ and $t \in (\frac{1}{2}, 1]$, if the following subordination holds:

$$\left(\frac{zf'(z)}{f(z)}\right)^\alpha \left(1 + \frac{zf''(z)}{f'(z)}\right)^{1-\alpha} \prec H(z, t) := \frac{1}{1 - 2tz + z^2} \quad (z \in D). \tag{2}$$

We note that if $t = \cos \theta$, $\theta \in (-\frac{\pi}{3}, \frac{\pi}{3})$, then

$$H(z, t) = \frac{1}{1 - 2tz + z^2}$$

$$= 1 + \sum_{n=1}^{\infty} \frac{\sin(n+1)\theta}{\sin \theta} z^n \quad (z \in D).$$

Thus,

$$H(z, t) = 1 + 2 \cos \theta z + (3 \cos^2 \theta - \sin^2 \theta)z^2 + \dots \quad (z \in D).$$

Following that, we can write

$$H(z, t) = 1 + U_1(t)z + U_2(t)z^2 + \dots \quad (z \in D, t \in (-1, 1)),$$

where $U_{n-1} = \frac{\sin(n \arccos t)}{\sqrt{1-t^2}}$ ($n \in \mathbb{N}$) are the Chebyshev polynomials of the second kind. Also, it is known that

$$U_n(t) = 2tU_{n-1}(t) - U_{n-2}(t),$$

and

$$U_1(t) = 2t,$$

$$U_2(t) = 4t^2 - 1,$$

$$U_3(t) = 8t^3 - 4t,$$

$$\vdots$$
(3)

The Chebyshev polynomials $T_n(t)$, $t \in [-1, 1]$, of the first kind have the generating function of the form

$$\sum_{n=0}^{\infty} T_n(t)z^n = \frac{1 - tz}{1 - 2tz + z^2} \quad (z \in D).$$

However, the Chebyshev polynomials of the first kind $T_n(t)$ and the second kind $U_n(t)$ are well connected by the following relationships:

$$\begin{aligned} \frac{dT_n(t)}{dt} &= nU_{n-1}(t), \\ T_n(t) &= U_n(t) - tU_{n-1}(t), \\ 2T_n(t) &= U_n(t) - U_{n-2}(t). \end{aligned}$$

In this paper, motivated by the earlier work of Dziok et al. [4], we use the Chebyshev polynomial expansions to provide estimates for the initial coefficients of univalent functions in $L(\alpha, t)$.

2. Coefficient bounds for the function class $L(\alpha, t)$

In this section, we propose to find the estimates on the Taylor–Maclaurin coefficients $|a_2|$ and $|a_3|$ for functions in the class $L(\alpha, t)$, which we introduced in Definition 1. We first state Theorem 2.

Theorem 2 *Let the function f given by (1) be in the class $L(\alpha, t)$. Then*

$$|a_2| \leq \frac{2t}{2 - \alpha}$$

and

$$|a_3| \leq \frac{(16 - 13\alpha + \alpha^2)t^2}{(3 - 2\alpha)(2 - \alpha)^2} + \frac{t}{3 - 2\alpha} - \frac{1}{2(3 - 2\alpha)}.$$

Proof Let $f \in L(\alpha, t)$. From (2), we have

$$\left(\frac{zf'(z)}{f(z)}\right)^\alpha \left(1 + \frac{zf''(z)}{f'(z)}\right)^{1-\alpha} = 1 + U_1(t)w(z) + U_2(t)w^2(z) + \dots, \tag{4}$$

for some analytic functions w such that $w(0) = 0$ and $|w(z)| < 1$, for all $z \in D$. From the equality (4), we obtain that

$$\left(\frac{zf'(z)}{f(z)}\right)^\alpha \left(1 + \frac{zf''(z)}{f'(z)}\right)^{1-\alpha} = 1 + U_1(t)c_1z + [U_1(t)c_2 + U_2(t)c_1^2]z^2 + \dots. \tag{5}$$

It is fairly well known that if $|w(z)| = |c_1z + c_2z^2 + c_3z^3 + \dots| < 1$, $z \in D$, then

$$|c_j| \leq 1, \quad \text{for all } j \in \mathbb{N}, \tag{6}$$

and

$$|c_2 - \mu c_1^2| \leq \max\{1, |\mu|\}, \quad \text{for } \mu \in \mathbb{R}. \tag{7}$$

It follows from (5) that

$$(2 - \alpha) a_2 = U_1(t)c_1, \tag{8}$$

$$2(3 - 2\alpha) a_3 + \frac{\alpha^2 + 5\alpha - 8}{2} a_2^2 = U_1(t)c_2 + U_2(t)c_1^2. \tag{9}$$

From (3) and (8), we obtain

$$|a_2| \leq \frac{2t}{2 - \alpha}. \tag{10}$$

Next, in order to find the bound on $|a_3|$, by using (8) in (9), we obtain

$$2(3 - 2\alpha) a_3 = U_1(t)c_2 + \left\{ U_2(t) - \frac{\alpha^2 + 5\alpha - 8}{2(2 - \alpha)^2} U_1^2(t) \right\} c_1^2. \tag{11}$$

Then, in view of (3) and (6), we have from (11)

$$|a_3| \leq \frac{(16 - 13\alpha + \alpha^2) t^2}{(3 - 2\alpha)(2 - \alpha)^2} + \frac{t}{3 - 2\alpha} - \frac{1}{2(3 - 2\alpha)}.$$

□

3. Fekete–Szegő inequalities for the function class $L(\alpha, t)$

In this section, we derive the Fekete–Szegő inequalities for functions in the class $L(\alpha, t)$, which is introduced by Definition 1. These inequalities are asserted by Theorem 3.

Theorem 3 *Let f given by (1) be in the class $L(\alpha, t)$. Then*

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{t}{3 - 2\alpha}; & \text{for } \mu \in [\mu_1, \mu_2] \\ \frac{t}{3 - 2\alpha} \left| \frac{4t^2 - 1}{2t} - \frac{\alpha^2 + 5\alpha - 8}{(2 - \alpha)^2} t - 4\mu \frac{(3 - 2\alpha)}{(2 - \alpha)^2} t \right|; & \text{for } \mu \notin [\mu_1, \mu_2] \end{cases},$$

where,

$$\mu_1 = \frac{2t^2(\alpha^2 - 13\alpha + 16) - (2 - \alpha)^2(1 + 2t)}{8(3 - 2\alpha)t^2}, \quad \mu_2 = \frac{2t^2(\alpha^2 - 13\alpha + 16) - (2 - \alpha)^2(1 - 2t)}{8(3 - 2\alpha)t^2}.$$

Proof From (8) and (11), we can easily see that

$$a_3 - \mu a_2^2 = \frac{U_1(t)}{2(3 - 2\alpha)} c_2 + \left\{ \frac{U_2(t)}{2(3 - 2\alpha)} - \frac{\alpha^2 + 5\alpha - 8}{4(3 - 2\alpha)(2 - \alpha)^2} U_1^2(t) - \mu \frac{U_1^2(t)}{(2 - \alpha)^2} \right\} c_1^2$$

and

$$|a_3 - \mu a_2^2| = \frac{U_1(t)}{2(3 - 2\alpha)} \left| c_2 + \left\{ \frac{U_2(t)}{U_1(t)} - \frac{\alpha^2 + 5\alpha - 8}{2(2 - \alpha)^2} U_1(t) - 2\mu \frac{(3 - 2\alpha) U_1(t)}{(2 - \alpha)^2} \right\} c_1^2 \right|.$$

Then, in view of (7), we conclude that

$$|a_3 - \mu a_2^2| \leq \frac{U_1(t)}{2(3-2\alpha)} \max \left\{ 1, \left| \frac{U_2(t)}{U_1(t)} - \frac{\alpha^2 + 5\alpha - 8}{2(2-\alpha)^2} U_1(t) - 2\mu \frac{(3-2\alpha)U_1(t)}{(2-\alpha)^2} \right| \right\}. \tag{12}$$

Finally, by using (3) in (12), we get

$$|a_3 - \mu a_2^2| \leq \frac{t}{3-2\alpha} \max \left\{ 1, \left| \frac{4t^2 - 1}{2t} - \frac{\alpha^2 + 5\alpha - 8}{(2-\alpha)^2} t - 4\mu \frac{(3-2\alpha)}{(2-\alpha)^2} t \right| \right\}.$$

Because $t > 0$, we have

$$\begin{aligned} & \left| \frac{4t^2 - 1}{2t} - \frac{\alpha^2 + 5\alpha - 8}{(2-\alpha)^2} t - 4\mu \frac{(3-2\alpha)}{(2-\alpha)^2} t \right| \leq 1 \\ \Leftrightarrow & \left\{ \frac{2t^2(\alpha^2 - 13\alpha + 16) - (2-\alpha)^2(1+2t)}{8(3-2\alpha)t^2} \leq \mu \leq \frac{2t^2(\alpha^2 - 13\alpha + 16) - (2-\alpha)^2(1-2t)}{8(3-2\alpha)t^2} \right\} \\ \Leftrightarrow & \mu_1 \leq \mu \leq \mu_2. \end{aligned}$$

□

In its special case when $\alpha = 0$, Theorem 3 leads us to Corollary 4.

Corollary 4 *If $f \in L(t)$, then*

$$\begin{aligned} & |a_2| \leq t; \\ & |a_3| \leq \frac{4t^2}{3} + \frac{t}{3} - \frac{1}{6}; \\ & |a_3 - \mu a_2^2| \leq \begin{cases} \frac{t}{3}; & \text{for } \mu \in [\mu_1, \mu_2] \\ \left| \frac{8t^2 - 1 - 6\mu t^2}{6} \right|; & \text{for } \mu \notin [\mu_1, \mu_2] \end{cases}. \end{aligned}$$

References

- [1] Altinkaya Ş, Yalçın S. Fekete-Szegő inequalities for certain classes of bi-univalent functions. International Scholarly Research Notices 2014; 2014: 327962.
- [2] Bulut S. Fekete-Szegő problem for subclasses of analytic functions defined by Komatu integral operator. Arab J Math 2013; 2: 177-183.
- [3] Doha EH. The first and second kind Chebyshev coefficients of the moments of the general-order derivative of an infinitely differentiable function. Int J Comput Math 1994; 51: 21-35.
- [4] Dziok J, Raina RK, Sokol J. Application of Chebyshev polynomials to classes of analytic functions. C R Acad Sci Paris Ser I 2015; 353: 433-438.
- [5] Fekete M, Szegő G. Eine Bemerkung über ungerade schlichte Funktionen. J Lond Math Soc 1933; 2: 85-89 (in German).
- [6] Mason JC. Chebyshev polynomials approximations for the L -membrane eigenvalue problem. SIAM J Appl Math 1967; 15: 172-186.

- [7] Orhan H, Magesh N, Balaji VK. Fekete-Szegő problem for certain classes of Ma-Minda bi-univalent functions. Afr Mat 2016; 27: 889-897.
- [8] Zaprawa Z. On Fekete-Szegő problem for classes of bi-univalent functions. Bull Belg Math Soc Simon Stevin 2014; 21: 169-178.