

Incidence of Type 1 Diabetes in Children Aged Below 18 Years during 2013-2015 in Northwest Turkey

© Sükran Poyrazoğlu¹, © Rüveyde Bundak^{1,2}, © Zehra Yavaş Abalı¹, © Hasan Önal³, © Sevil Sarıkaya³, © Abdurrahman Akgün³, © Serpil Baş⁴, © Saygın Abalı^{4,5}, © Abdullah Bereket⁴, © Erdal Eren⁶, © Ömer Tarım⁶, © Ayla Güven^{7,8}, © Metin Yıldız⁸, © Derya Karaman Aksakal⁸, © Ayşegül Yüksel⁹, © Gülcan Seymen Karabulut⁹, © Şükrü Hatun^{9,10}, © Tolga Özgen¹¹, © Yaşar Cesur¹¹, © Mehmet Azizoğlu¹², © Emine Dilek¹², © Filiz Tütüncüler¹², © Esra Papatya Çakır^{13,14}, © Bahar Özcabı^{15,16}, © Olcay Evliyaoğlu¹⁵, © Songül Karadeniz¹⁵, © Fatma Dursun¹⁷, © Semih Bolu¹⁸, © İlknur Arslanoğlu¹⁸, © Gül Yeşiltepe Mutlu^{9,16}, © Heves Kırmızıbekmez¹⁶, © Pınar İşgüven¹⁹, © Ala Üstyol²⁰, © Erdal Adal²¹, © Ahmet Uçar²², © Nurcan Cebeci²³, © Didem Bezen²⁴, © Çiğdem Binay²⁵, © Serap Semiz²⁶, © Hüseyin Anıl Korkmaz²⁷, © Nihal Memioğlu²⁸, © Elif Sağsak²⁹, © Havva Nur Peltek³⁰, © Melek Yıldız³, © Teoman Akçay³¹, © Serap Turan⁴, © Tülay Güran⁴, © Zeynep Atay^{4,21}, © Neşe Akcan^{8,32}, © Filiz Çizmecioglu⁹, © Oya Ercan¹⁵, © Aydılek Dağdeviren¹⁵, © Firdevs Baş¹, © Halim İşsever³³, © Feyza Darendeliler¹

¹Istanbul University Istanbul Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

²University of Kyrenia Faculty of Medicine, Department of Pediatric Endocrinology, Kyrenia, Turkish Republic of North Cyprus

³Kanuni Sultan Süleyman Training and Research Hospital, Clinic of Pediatric Endocrinology and Metabolic Disease, Istanbul, Turkey

⁴Marmara University Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

⁵Kartal Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

⁶Uludağ University Faculty of Medicine, Department of Pediatric Endocrinology, Bursa, Turkey

⁷Amasya University Faculty of Medicine, Department of Pediatric Endocrinology, Amasya, Turkey

⁸Göztepe Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

⁹Kocaeli University Faculty of Medicine, Department of Pediatric Endocrinology, Kocaeli, Turkey

¹⁰Koç University Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

¹¹Bezmialem Vakıf University Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

¹²Trakya University Faculty of Medicine, Department of Pediatric Endocrinology, Edirne, Turkey

¹³Bakırköy Dr. Sadi Konuk Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

¹⁴Şevket Yılmaz Training and Research Hospital, Clinic of Pediatric Endocrinology, Bursa, Turkey

¹⁵Istanbul University Cerrahpaşa Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

¹⁶Zeynep Kamil Women's and Children's Disease Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

¹⁷Ümraniye Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

¹⁸Düzce University Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

¹⁹Sakarya University Faculty of Medicine, Department of Pediatric Endocrinology, Sakarya, Turkey

²⁰Haseki Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

²¹Medipol University Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey

²²Şişli Etfal Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

²³Derince Training and Research Hospital, Clinic of Pediatric Endocrinology, Kocaeli, Turkey

²⁴Okmeydanı Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

²⁵Çorlu State Hospital, Clinic of Pediatric Endocrinology, Tekirdağ, Turkey

²⁶Acıbadem University Faculty of Medicine, Clinic of Pediatric Endocrinology, Istanbul, Turkey

²⁷Balıkesir Atatürk State Hospital, Clinic of Pediatric Endocrinology, Balıkesir, Turkey

²⁸American Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

²⁹Gaziosmanpaşa Taksim Training and Research Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

³⁰Edirne Sultan 1. Murat State Hospital, Clinic of Pediatric Endocrinology, Edirne, Turkey

³¹Medical Park Gaziosmanpaşa Hospital, Clinic of Pediatric Endocrinology, Istanbul, Turkey

³²University of Near East Faculty of Medicine, Department of Pediatric Endocrinology, Nicosia, Turkish Republic of North Cyprus

³³Istanbul University Istanbul Faculty of Medicine, Department of Public Health, Istanbul, Turkey



Address for Correspondence: Sükran Poyrazoğlu MD, Istanbul University Istanbul Faculty of Medicine, Department of Pediatric Endocrinology, Istanbul, Turkey
Phone: +90 212 414 20 00 **E-mail:** sukranpoyrazoglu@yahoo.com **ORCID ID:** orcid.org/0000-0001-6806-9678

©Copyright 2018 by Turkish Pediatric Endocrinology and Diabetes Society

The Journal of Clinical Research in Pediatric Endocrinology published by Galenos Publishing House.

Conflict of interest: None declared

Received: 22.01.2018

Accepted: 22.05.2018

What is already known on this topic?

Incidence of type 1 diabetes mellitus (T1DM) peaked in the age groups 5-9 and 10-14 years. Diagnosis of T1DM showed a seasonal pattern peaking in autumn-winter.

What this study adds?

The incidence of type 1 diabetes mellitus in children and adolescents aged 0-17 years was 8.99/100,000 during 2013-2015 in the Northwestern region of Turkey and constant over the course of these 3 years.

Abstract

Objective: To assess the incidence of type 1 diabetes mellitus (T1DM) in children under 18 years of age in the northwest region of Turkey during 2013-2015.

Methods: All newly diagnosed T1DM cases were recorded prospectively during 2013-2015. Total, as well as gender and age group specific (0-4, 5-9, 10-14 and 15-17 age) mean incidences per 100,000 per year were calculated.

Results: There were 1,773 patients diagnosed during 2013-2015 (588 cases in 2013, 592 cases in 2014, 593 cases in 2015). Of these, 862 (48.6%) were girls and 911 (51.4%) were boys. The mean age at diagnosis was 9.2 ± 4.2 years and it was not significantly different between girls (9.0 ± 4.1 years) and boys (9.4 ± 4.4 years) ($p = 0.052$). The crude mean incidence was 8.99/100,000 confidence interval (CI) (95% CI: 8.58-9.42). Although mean incidence was similar between boys [8.98/100,000 (CI: 8.40 to 9.58)] and girls [9.01/100,000 (CI: 8.42 to 9.63)], there was male predominance in all groups except for 5-9 year age group. The standardized mean incidence was 9.02/100,000 according to the World Health Organization standard population. The mean incidence for the 0-4, 5-9, 10-14 and 15-17 age groups was 6.13, 11.68, 11.7 and 5.04/100,000 respectively. The incidence of T1DM was similar over the course of three years ($p = 0.95$). A significant increase in the proportion of cases diagnosed was observed in the autumn-winter seasons.

Conclusion: The northwest region of Turkey experienced an intermediate incidence of T1DM over the period of the study.

Keywords: Type 1 diabetes mellitus, childhood, incidence

Introduction

Type 1 diabetes mellitus (T1DM) is a common, chronic disease in children and adolescents. In many populations, an increase in the incidence of T1DM in children has been observed (1,2,3). Studies have shown that the incidence of T1DM varies widely between and within countries (1,2,3,4). Seasonal variations in the presentation and gender differences in incidence of T1DM have been reported (5,6,7,8).

In Turkey, data on incidence and incidence trends of childhood T1DM are limited. Our aim was to determine the incidence of T1DM in children and adolescents (aged under 18 years) during the years 2013-2015 in the northwestern region of Turkey and to analyze the seasonal presentation pattern of T1DM in these children.

Methods

Turkey is divided into seven geographical regions determined by topography and climate and defined by central government. The northwestern region, where this prospective study was conducted, is one of these regions. All children younger than 18 years of age, diagnosed as T1DM in pediatric endocrinology units in this region during 2013-2015 were included in the study. The pediatric

endocrinology units' locations were 11 university hospitals, 15 state hospitals and one private hospital.

The diagnosis of T1DM was made by the pediatric endocrinologist who took care of the child, according to the accepted criteria of the International Society for Pediatric and Adolescent Diabetes (9). The date of diagnosis of diabetes was accepted as the day of the first insulin injection.

In the Turkish health care system, all children aged 0-17 years with T1DM are referred to a pediatric endocrinology department for treatment. Over the three year period (2013-2015), data on all hospitalized or referred new cases in the institutions in the Northwestern region of the country were reported to our team on a special form containing information about the patient's personal identification number, sex, date of birth, date of diagnosis and some clinical and laboratory data. All forms were sent monthly to one investigator (SP) for data collection and verification. We excluded children with type 2 diabetes mellitus, neonatal diabetes, maturity onset diabetes of youth, transient hyperglycemia, and diabetes caused by other conditions (chemotherapy, cystic fibrosis, etc).

Statistical Analysis

Incidence of T1DM was calculated using the numbers of patients reported for each year by age (0-4, 5-9, 10-14

and 15-17 years aged) and gender groups (girls and boys). Annual numbers for the age groups in the geographical area were used as denominators, and incidence (per 100,000 per year) was calculated with 95% CIs, assuming a Poisson distribution. The annual population sizes were obtained from the Turkish census data of 2013-2015 from the address-based population registration system of the Turkish Statistical Institute. For comparison with data from other countries, the incidence was standardized by the direct method according to the age distribution of the world population (10).

The percentage of patients diagnosed during each calendar month was calculated in age groups for both sexes and then compared, to identify any seasonal variation in diagnosis of T1DM.

In order to assess the significance of the differences between the groups, normality of variables was tested by Kolmogorov Smirnov test; Mann-Whitney U and chi-square tests were used. Results are reported as means \pm SD. Two-tailed p values were calculated. Statistical significance was accepted as $p < 0.05$.

Results

A total of 1773 cases were identified over the three year period (588 cases in 2013, 592 cases in 2014, 593 cases in 2015). Of these, 862 (48.6%) were girls and 911 (51.4%) were boys, giving a male to female ratio of 1.05:1. The mean age at diagnosis was 9.2 ± 4.2 years and showed no sex difference (9.0 ± 4.1 years in the girls and 9.4 ± 4.4 years in the boys, $p = 0.052$). Table 1 shows mean ages and distribution of the patients by age groups over the three year period. The proportion of newly diagnosed T1DM cases was highest among children aged 5-9 years (35.9%), followed by

the age groups 10-14 years (35.3%), 0-4 years (19.1%) and 15-17 years (9.6%).

The crude mean annual incidence in children aged 0-17 years over this period was 8.99 per 100,000 [95% confidence interval (CI): 8.58 to 9.42]. The standardized mean incidence was 9.02 per 100,000 according to the World Health Organization (WHO) standard population.

There was no significant difference between the mean annual incidence figures for boys [8.98/100.000 (CI: 8.40 to 9.58)] and girls [9.01/100.000 (CI: 8.42 to 9.63)] during the study period ($p = 0.95$) (Table 2). The mean annual incidence for the 0-4 year age group was 6.13/100.000. Incidence increased significantly with age, reaching a peak in the age groups 5-9 and 10-14 years. It was 11.68/100.000 for the 5-9 year age group and 11.7/100.000 for the 10-14 year age group and subsequently the incidence declined at age 15-17 years. The lowest incidence was seen in the age group 15-17 years (5.04/100,000) (Table 2). The incidence of the age group 0-14 years was 9.82/100,000 (95% CI: 9.34 to 10.31). Male predominance was seen in all groups except for the 5-9 years age group (Table 2). The incidence of T1DM was similar over the course of the three years (Table 2, $p = 0.95$).

A significant increase in proportion of diagnosis of T1DM was observed in the autumn-winter seasons (Figure 1). It was similar over the three year period in all age and gender groups.

Discussion

In this study we investigated the incidence of T1DM in children residing in the Northwestern region of Turkey, our results demonstrate that the incidence of T1DM here is

Table 1. Mean ages at diagnosis and distribution by age groups of newly diagnosed type 1 diabetes mellitus cases over the three year period

	2013	2014	2015	2013-2015
Total number of cases	588	592	593	1773
Boys	279	309	323	911
Girls	309	283	270	862
Mean age at diagnosis (years)				
Total group	9.2 ± 4.3	8.9 ± 4.3	9.5 ± 4.1	9.2 ± 4.2
Boys	9.6 ± 4.5	9.1 ± 4.4	9.5 ± 4.3	9.4 ± 4.4
Girls	8.9 ± 4.1	8.7 ± 4.2	9.4 ± 3.9	9.0 ± 4.1
Proportion by age groups (%)				
0-4 years	20	20.9	16	19.1
5-9 years	32.3	38.2	37	35.9
10-14 years	39.1	31.4	35.4	35.3
15-17 years	8.3	9.5	11.1	9.6

intermediate (8.99/100,000) in the pediatric age group (4). In Turkey, there are only a few reports on the epidemiology of T1DM in children and most of them focused on children below 15 years of age (11,12,13,14). Our study is one of the few population-based reports presenting the incidence of T1DM among children in Turkey.

Considerable differences in incidence rates for T1DM have been reported from different countries, and even within the same country (1,2,3,4). Recently, in 2013, a nationwide incidence of T1DM among Turkish children was reported and this study covered T1DM incidence in Turkey divided 5 geographic regions (14) with notable differences in

incidence across the five regions. The northwestern region is a developing part of Turkey and there have been big changes in the economy, urbanization and lifestyles in recent decades in this region. A slightly higher incidence (10.1/100,000 per year) was reported in the all western part of Turkey in this nationwide study and the results were consistent with our data (9.01/100,000 per year). However, the methodology for case ascertainment used in our study is different from the nationwide study. The nationwide study used data from the universal health insurance system about prescriptions for essential medicines for diabetics for the calculation of incidence. In our study the data were collected prospectively from T1DM patient data from pediatric endocrinology units in the region.

We cannot detect the incidence trends from our study due to the short period covered and the lack of epidemiological data before our study in children younger than 18 years in Turkey. Our neighbour countries reported intermediate rates for incidence, similar to our results (2,3,15) and reported an increase in incidence of T1DM over time. Although our observation period was very short, incidence was quite stable over the three year period. Although the global increase in the incidence of T1DM is widely recognized in recent decades, some studies in populations with a higher incidence of T1DM have demonstrated that the increase in the incidence slowed down in the last decade (16,17,18). In the EuroDiab study, it was reported that between 2001 and 2009 the




Figure 1. Distribution of age of onset of type 1 diabetes according to months of the year

Table 2. Incidence of type 1 diabetes mellitus over the three year period

Incidence (95% CI)	2013-2015	2013	2014	2015	p
Total group	8.99	9.01	8.99	8.98	0.95
Boys	8.98	8.31	9.11	9.51	0.07
Girls	9.01	9.76	8.85	8.43	0.1
By age groups (total group)					
0-4 years	6.13	6.6	6.74	5.09	0.92
5-9 years	11.68	10.56	12.37	12.10	0.85
10-14 years	11.7	12.78	10.42	11.9	0.48
15-17 years	5.04	4.37	4.93	5.82	0.60
Boys					
0-4 years	6.39	6.26	6.98	5.94	0.78
5-9 years	10.26	8.43	11.08	11.25	0.06
10-14 years	12.73	12.29	12.39	13.51	0.41
15-17 years	5.19	5.01	4.26	6.31	0.32
Girls					
0-4 years	5.86	6.96	6.48	4.18	0.07
5-9 years	13.18	12.8	13.74	13.01	0.91
10-14 years	10.6	13.31	8.32	10.18	0.06
15-17 years	4.88	3.69	5.65	5.28	0.23

CI: confidence interval

increase in T1DM incidence was significantly different in regions within Europe and the highest increase occurred in Central Eastern European countries while Finland, other Nordic countries and the Czech Republic showed a much lower increase or a stabilization in the incidence of T1DM (16,17,19,20,21). An average relative increase of 3-4% per year has been reported worldwide (22). Environmental factors are thought as the most likely reason for this increase in incidence (3,20,23). For this reason, evaluation of incidence in different regions is important and warranted.

In our cohort, although the mean annual incidence for boys and girls was similar, a male predominance was seen in all age groups except for the age group 5-9 years. The female predominance in the 5-9 years age group could be due to the earlier onset of puberty in girls than in boys. Gender differences in T1DM have been identified in many studies (4,6,24,25,26,27). Overall, high incidence countries tend to have a slight male predominance and low incidence countries a female predominance (4,24,25,26,27). Karvonen et al (4) found that 88% of low incidence populations were predominantly girls and patients in high incidence populations were more likely to be predominantly boys. In Sardinia, a very high incidence area, a male predominance is reported in the 0-14 year age group (26). The Danish Study Group of Diabetes in Childhood reported male predominance in their population (24). However, no significant difference in T1DM incidence between boys and girls was observed in Shanghai or in Kuwait (25,27).

Age differences in T1DM incidence have also been observed in previous studies (2,15,27,28,29). The incidence in our cohort increased with age in both sexes and was highest in the 5-14 year age group. This was followed by a decrease in the 15-17 year age group. Incidence was similar in children aged 5-9 and 10-14 years in our study. The youngest age group (0-4 years) had a lower incidence as compared with older children (5-14 years). This difference in incidence by age groups has also been shown in other counties. The WHO analyzed standardized incidence data on T1DM in the Multinational Project (DIAMOND) in 112 centres from 57 countries during 1990-1999. The DIAMOND study showed that 5-9 years old children had a higher risk of developing T1DM compared with 0-4 year old children (2). Some countries reported a high incidence in the 5-9 years old group, but others found the highest incidence in children aged 10-14 years (2,15,27,28,29).

Our cohort showed a significant seasonal variation in diagnosis of T1DM. More cases were diagnosed during

autumn and winter months, which are the cooler seasons in the Northwest region. This seasonality of diagnosis of T1DM was identified in both sexes and in all age groups and thus seems to be a robust finding. Although Turkey is situated in the Mediterranean geographical location, the diverse regions have different climates because of irregular topography. To evaluate the impact of weather on incidence of T1DM in Turkey, each region should be evaluated separately. Similar to our results some countries show significant seasonality in diagnosis for all age groups, with higher incidence rates in the winter period (5,7,8). Some countries reported no seasonality in all age groups or absent in some age groups (5,8,30,31,32). Different interpretations have been suggested for this seasonal pattern in diagnosis of T1DM, including physical activity, stress, viral infections and vitamin D synthesis during different seasons (5,7,8,33). The DIAMOND group reported seasonality in T1DM incidence with winter or summer peaks in 40% of all participating centers depending on the geographic position of the country (2). It was shown that in Japan there was a bimodal pattern in the diagnosis of T1DM, that is common in April/May and in December with no seasonal pattern of incidence among preschool children (6).

Study Limitation

The limitation of our study was the short duration of the registry. It would be important to continue monitoring incidence of T1DM in the same region and evaluate trends.

Conclusion

To conclude, this is the first paper that analyzes the time-related trends in the incidence of T1DM in Turkish children aged from 0 to 17 years in the Northwest region of Turkey. The results showed an intermediate incidence of T1DM and a similar mean annual incidence between boys and girls. Considering the increasing incidence of T1DM worldwide, we suggest that it would be important to follow trends in incidence in the next few years in this same region to determine the possible triggering factors and also to develop preventive strategies.

Ethics

Ethics Committee Approval: This study was approved by the Local Ethical Committee of İstanbul Faculty of Medicine.

Informed Consent: Informed consent was obtained from the families and children.

Peer-review: Externally and internally peer-reviewed.

Authorship Contributions

Concept: Şükran Poyrazoğlu, Rüveyde Bundak, Design: Şükran Poyrazoğlu, Rüveyde Bundak, Data Collection and Processing: All authors, Analysis and Interpretations: Şükran Poyrazoğlu, Rüveyde Bundak, Halim İşsever, Writing: Şükran Poyrazoğlu, Rüveyde Bundak, Feyza Darendeliler.

Financial Disclosure: The authors declare that this study received no financial support.

References

1. Patterson C, Guariguata L, Dahlquist G, Soltész G, Ogle G, Silink M. Diabetes in the young-a global view and worldwide estimates of numbers of children with type 1 diabetes. *Diabetes Res Clin Pract* 2014;103:161-175.
2. DIAMOND Project Group. Incidence and trends of childhood type 1 diabetes worldwide 1990-1999. *Diabet Med* 2006;23:857-866.
3. Patterson CC, Gyürüs E, Rosenbauer J, Cinek O, Neu A, Schober E, Parslow RC, Joner G, Svensson J, Castell C, Bingley PJ, Schoenle E, Jarosz-Chobot P, Urbonaitė B, Rothe U, Krzysnik C, Ionescu-Tirgoviste C, Weets I, Kocova M, Stipančić G, Samardžić M, de Beaufort CE, Green A, Dahlquist GG, Soltész G. Trends in childhood type 1 diabetes incidence in Europe during 1989-2008: evidence of non-uniformity over time in rates of increase. *Diabetologia* 2012;55:2142-2147.
4. Karvonen M, Viik-Kajander M, Moltchanova E, Libman I, Laporte R, Tuomilehto J. The Diabetes Mondiale (Diamond) Project Group. Incidence of childhood type 1 diabetes worldwide. *Diabetes Care* 2000;23:1516-1526.
5. Patterson CC, Gyürüs E, Rosenbauer J, Cinek O, Neu A, Schober E, Parslow RC, Joner G, Svensson J, Castell C, Bingley PJ, Schoenle E, Jarosz-Chobot P, Urbonaitė B, Rothe U, Krzysnik C, Ionescu-Tirgoviste C, Weets I, Kocova M, Stipančić G, Samardžić M, de Beaufort CE, Green A, Soltész G, Dahlquist GG. Seasonal variation in month of diagnosis in children with type 1 diabetes registered in 23 European centers during 1989-2008: little short-term influence of sunshine hours or average temperature. *Pediatr Diabetes* 2015;16:573-580.
6. Onda Y, Sugihara S, Ogata T, Yokoya S, Yokoyama T, Tajima N. Type 1 Diabetes (T1D) Study Group. Incidence and prevalence of childhood-onset Type 1 diabetes in Japan: the T1D study. *Diabet Med* 2017;34:909-915.
7. Svensson J, Lyngaae-Jørgensen A, Carstensen B, Simonsen LB, Mortensen HB. Danish Childhood Diabetes Registry. Long-term trends in the incidence of type 1 diabetes in Denmark: the seasonal variation changes over time. *Pediatr Diabetes* 2009;10:248-254.
8. Samuelsson U, Carstensen J, Löfman O, Nordfeldt S. Seasonal variation in the diagnosis of type 1 diabetes in south-east Sweden. *Diabetes Res Clin Pract* 2007;76:75-81.
9. Craig M, Hattersley A, Donaghue K. International Society For Pediatric And Adolescent Diabetes. ISPAD Clinical Practice Consensus Guidelines 2006-2007. Definition, Epidemiology And Classification. *Pediatr Diabetes* 2006;7:343-351.
10. Ahmad OB, Boschi-Pinto C, Lopez AD, Murray C, Lozano R, Inoue M. Age Standardization of Rates: A New Who Standard. EIP/GPE/EBD. World Health Organization, 2001.
11. Akesen E, Turan S, Guran T, Atay Z, Save D, Bereket A. Prevalence of type 1 diabetes mellitus in 6-18-yr-old school children living in Istanbul, Turkey. *Pediatr Diabetes* 2011;12:567-571.
12. Demirbilek H, Ozbek MN. Prevalence of type 1 diabetes mellitus in schoolchildren 6-18 years old in Diyarbakır, Southeastern Anatolian Region of Turkey. *Turk J Med Sci* 2013;43:768-774.
13. Demirbilek H, Ozbek MN, Baran RT. Incidence of type 1 diabetes mellitus in Turkish children from the southeastern region of the country: a regional report. *J Clin Res Pediatr Endocrinol* 2013;5:98-103.
14. Yeşilkaya E, Cinaz P, Andıran N, Bideci A, Hatun Ş, Sarı E, Türker T, Akgül Ö, Saldır M, Kılıçaslan H, Açıkcel C, Craig ME. First report on the nationwide incidence and prevalence of Type 1 diabetes among children in Turkey. *Diabet Med* 2017;34:405-410.
15. Serban V, Brink S, Timar B, Sima A, Vlad M, Timar R, Vlad A. An increasing incidence of type 1 diabetes mellitus in Romanian children aged 0 to 17 years. *J Pediatr Endocrinol Metab* 2015;28:293-298.
16. Harjutsalo V, Sund R, Knip M, Groop PH. Incidence of type 1 diabetes in Finland. *JAMA* 2013;310:427-428.
17. Skrivarhaug T, Stene LC, Drivvoll AK, Strom H, Joner G. Incidence of type 1 diabetes in Norway among children aged 0-14 years between 1989 and 2012: has the incidence stopped rising? Results from the Norwegian Childhood Diabetes Registry. *Diabetologia* 2014;57:57-62.
18. Berhan Y, Waernbaum I, Lind T, Möllsten A, Dahlquist G. Thirty years of prospective nationwide incidence of childhood type 1 diabetes: the accelerating increase by time tends to level off in Sweden. *Diabetes* 2011;60:577-581.
19. Soltész G, Patterson CC, Dahlquist G, EURODIAB Study Group. Worldwide childhood type 1 diabetes incidence-what can we learn from epidemiology? *Pediatr Diabetes* 2007;8:6-14.
20. Patterson CC, Dahlquist G, Gyürüs E, Green A, Soltész G, EURODIAB Study Group. Incidence trends for childhood type 1 diabetes in Europe during 1989-2003 and predicted new cases 2005-20: a multicentre prospective registration study. *Lancet* 2009;373:2027-2033.
21. Cinek O, Kulich M, Sumnik Z. The incidence of type 1 diabetes in Young Czech children stopped rising. *Pediatr Diabetes* 2012;13:559-563.
22. Tuomilehto J. The emerging global epidemic of type 1 diabetes. *Curr Diab Rep* 2013;13:795-804.
23. Bodin J, Stene LC, Nygaard UC. Can exposure to environmental chemicals increase the risk of diabetes Type 1 development? *BioMed Res Int* 2015; 2015:208947.
24. Svensson J, Carstensen B, Mortensen HB, Borch-Johnsen K. Danish Study Group of Childhood Diabetes: Early childhood risk factors associated with type 1 diabetes-is gender important? *Eur J Epidemiol* 2005;20:429-434.
25. Zhao Z, Sun C, Wang C, Li P, Wang W, Ye J, Gu X, Wang X, Shen S, Zhi D, Lu Z, Ye R, Cheng R, Xi L, Li X, Zheng Z, Zhang M, Luo F. Rapidly rising incidence of childhood type 1 diabetes in Chinese population: epidemiology in Shanghai during 1997-2011. *Acta Diabetol* 2014;51:947-953.
26. Casu A, Pascutto C, Bernardinelli L, Songini M. Type 1 diabetes among Sardinian children is increasing. The Sardinian diabetes register for children aged 0-14 years (1989-1999). *Diabetes Care* 2004;27:1623-1629.
27. Shaltout AA, Wake D, Thanaraj TA, Omar DM, Al-AbdulRazzaq D, Channanath A, AlKandari H, Abdulrasoul M, Miller S, Conway N, Tuomilehto J, Davidsson L; Steering Group for the Study of Childhood Diabetes in Kuwait. Incidence of type 1 diabetes has doubled in Kuwaiti children 0-14 years over the last 20 years. *Pediatr Diabetes* 2017;18:761-766.
28. Wu HB, Zhong JM, Hu RY, Wang H, Gong WW, Pan J, Fei FR, Wang M, Guo LH, Yang L, Yu M. Rapidly rising incidence of Type 1 diabetes in

- children and adolescents aged 0-19 years in Zhejiang, China, 2007 to 2013. *Diabet Med* 2016 ;33:1339-1346.
29. Fernandez-Ramos C, Arana-Arri E, Jimenez-Huertas P, Vela A, Rica I. Incidence of childhood-onset type 1 diabetes in Biscay, Spain, 1990-2013. *Pediatr Diabetes* 2017;18:71-76.
30. Ye J, Chen RG, Ashkenazi I, Laron Z. Lack of seasonality in the month of onset of childhood IDDM (0.7-15 years) in Shanghai, China. *J Pediatr Endocrinol Metab* 1998;11:461-464.
31. Karvonen M, Tuomilehto J, Virtala E, Pitkaniemi J, Reunanen A, Tuomilehto-Wolf E, Akerblom HK. Seasonality in the clinical onset of insulin-dependent diabetes mellitus in Finnish children. Childhood Diabetes in Finland (DiMe) Study Group. *Am J Epidemiol* 1996 15;143:167-176.
32. Padaiga Z, Tuomilehto J, Karvonen M, Dahlquist G, Podar T, Adojaan B, Urbonaite B, Zalinkevicius R, Brigis G, Virtala E, Kohtamäki K, Cepaitis Z, Tuomilehto-Wolf E. Seasonal variation in the incidence of Type 1 diabetes mellitus during 1983 to 1992 in the countries around the Baltic Sea. *Diabet Med* 1999;16:736-743.
33. Sloka S, Grant M, Newhook LA. Time series analysis of ultraviolet B radiation and type 1 diabetes in Newfoundland. *Pediatr Diabetes* 2008;9:81-86.